



Gruppo Umbro Mineralogico Paleontologico

FOSSILS & MINERALS

Review



SPECIAL ISSUE

WORLD CATALOG OF FOSSIL CANTHARIDAE

N° 2/2017

WORLD CATALOG OF FOSSIL CANTHARIDAE

Fabrizio FANTI
fantiff@libero.it

Abstract

A catalogue and bibliography of the world fossil Cantharidae are given. *Rhagonycha germari* (Heer, 1847) **n. comb.**, *Rhagonycha tertaria* (Heer, 1847) **n. comb.** and *Rhagonycha hesperus* (Wickham, 1914) **n. comb.** are transferred from *Telephorus* Schaeffer, 1766 = *Cantharis* Linnaeus, 1758 to *Rhagonycha* Eschscholtz, 1830. *Atalantycha humata* (Wickham, 1913) **n. comb.** is transferred from *Telephorus* Schaeffer, 1766 = *Cantharis* Linnaeus, 1758 to *Atalantycha* Kazantsev, 2005. *Lycocerus guttula* (J. Zhang, 1989) **n. comb.** is transferred from *Cantharis* Linnaeus, 1758 to *Lycocerus* Gorham, 1889. *Curticantharis* J. Zhang, 1989 = *Themus* Motschulsky, 1858 **n. syn.** *Themus capacis* (J. Zhang, 1989) **n. comb.**, *Themus thermophilus* (J. Zhang, 1989) **n. comb.**, *Themus trapezialis* (J. Zhang, Sun & X. Zhang, 1994) **n. comb.** and *Themus* sp. (J. Zhang, 1989) **n. comb.** are transferred from *Curticantharis* J. Zhang, 1989 to *Themus* Motschulsky, 1858. *Telephorus tertarius oeningensis* Heer, 1847 **n. syn.** and *Telephorus tertarius radobojanus* Heer, 1847 **n. syn.** are synonymized with *Rhagonycha tertaria* (Heer, 1847). Lectotype and paralectotype of *Rhagonycha tertaria* (Heer, 1847) are designated. *Malthacus deceptus* (W. J. Brown, 1940) **n. comb.** is the correct combination of *Podabrus deceptus* Brown = *Dichelotarsus deceptus*. A preliminary list of subfossil Cantharidae is also provided. Furthermore the Coleoptera Lampyridae *Lamprohiza fossilis* (Beier, 1952) **n. comb.** is transferred from *Phausis* LeConte, 1851 to *Lamprohiza* Motschulsky, 1853.

Key words: Catalog, worldwide, Coleoptera Cantharidae, paleontology

1. Introduction

The family Cantharidae is quite frequently found in fossil record, but currently there are only old or incomplete catalogs (Handlirsch 1906-1908; Spahr 1981a [bibliography], 1981b; Carpenter 1992), and only recently, on some lists and websites, are cited more species (Handlirsch 1906-1908; Spahr 1981a [bibliography], 1981b; Carpenter 1992), or we find various data for all fossil species (The Paleobiology Database). Also the world catalog (Delkeskamp 1939, 1977) does not mention these species, therefore with this article updated on 31 December 2016, I am finding and listing all the fossil species, including all citations that I know (some may be missing), and provide a preliminary list of the subfossil (Holocene) species (Table 2.); furthermore furnished an appendix (Appendix 1.) with the etymology and type species for the genera.

Acknowledgements

I am grateful to: Susana Adamonis and Cecilia Soledad Cataldo (Buenos Aires University), Vitalii I. Alekseev, Clive A. Auton, Gabriel Biffi (Brazil), Albertas Bitinas (Klaipėda University - Lithuania), Guido Bonamie (Merendree - Belgium), Jan Bosselaers, Christopher J. Duffin (Natural History Museum, London), Scott A. Elias, Michael S. Engel, Yun Hsiao (Taiwan); Jin Zhenyu (Sun Yat-Sen University, Guangzhou - China), Sergey V. Kazantsev; Wiesław Krzemieński (Museum PAN, Krakow), Ulrich Kotthoff (CeNak - Hamburg University), Jan-Michael Lange (Senckenberg Research Institute), Carlo Maria Legittimo, Cosmin-Ovidiu Manci (Romania), Yūichi Okushima (Kurashiki Museum of Natural History), Weston Opitz, Maurizio Pavesi, Evgeny E. Perkovsky, George O. Poinar Jr., Ren Dong and Wang Yongjie (College of Life Sciences, Capital Normal University - Beijing, China), Mónica Morayma Solórzano Kraemer, Francesco Vitali (Luxembourg), the Biodiversity Heritage Library (archive.org), the George Peabody Library (Paul Espinosa), the Naturalis Biodiversity Center - Netherlands Entomological Society, the Sutro Library - California State Library, and the Biblioteca Civica Gianfranco Contini of Domodossola for help in finding some papers.

And gratefully extend my thanks to: Roberto Antonio Pantaleoni and Francesco Vitali for the councils in some systematic problems, and Anders L. Damgaard, Sieghard Ellenberger, Artur R. Michalski, George O. Poinar for the courtesy of the photographs.

I wish to dedicate this article to the memory of Milton Sanderson (entomologist at Illinois Natural History Survey) and express my hearty thanks to his son Steven Sanderson for the English revision.

2. Methods

All named fossil taxa of the family Cantharidae are listed according to their current classification, and this is based on the articles of Ramsdale (2002), Kazantsev & Brancucci (2007) and Fanti (2014). The classification for the species is in the systematic-alphabetical order, and for the undescribed specimens also for the date of citation. For each species are indicated: 1- valid name and author(s) and date of description; 2- the original name combination and page(s) of description (excluded names in index); 3- eventually incorrect spelling or unjustified emendations; 4- all bibliographic references (although it is only mentioned the author or at least is evident the reference to one or more species) with number of pages, but for these are excluded species

names in index; 5- type horizon; 6- type locality; 7- preservation and lithology; 8- collection where is deposited the holotype and, eventually, the paratypes or other specimens, with number accession; 9- comments and notes, particularly focused on sex, body length, coloration, relationship with extant or fossil species, syninclusions and others deemed interesting news.

3.Systematic

Family Cantharidae Imhoff, 1856 (1815)

Subfamily Cantharinae Imhoff, 1856 (1815)

Tribe Cantharini Imhoff, 1856 (1815)

Genus *Atalantycha* Kazantsev, 2005

†***Atalantycha humata*** (Wickham, 1913) [*n. comb.*]

Telephorus humatus Wickham, 1913b: 360, 362, 366 (explanation of plates), Plate XXXVIII. Fig. 2. (but indicated as Plate I. Fig. 2. in the text and explanation) [under family Lampyridae] Lucas 1915: 135; International Catalogue of Scientific Literature 1916a: 95; Meunier 1920b: 159; Wickham 1920: 353 [as *Cantharis humatus*]; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis humata*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database [as *Cantharis humatus*].

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/

adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale, volcanioclastic.

COLLECTIONS. Museum of Princeton University, No. 5984.

COMMENTS AND NOTES. Body length: 7.50 mm. Very similar in form and size to the recent *Telephorus bilineatus* (= *Atalantycha bilineata* (Say) (Wickham 1913b: 362), and for this, and for the original drawing with pronotum rounded at sides and angles, second antennomer short and particularly the habitus short but stocky, I tentatively transferred this species to genus *Atalantycha* Kazantsev. Also for Carpenter (1992: 331), the assignment to genus *Cantharis* was doubt. The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

Genus *Cantharis* Linnaeus, 1758

Subgenus (*Cantharis*) Linnaeus, 1758

†***Cantharis (Cantharis) attavina*** (Heer, 1847)*¹

Telephorus attavinus Heer, 1847: 149-150, 225, Taf. IV. Fig. 15., Fig. 15. b.

Telephorus atavinus Pictet, 1854: 334 (incorrect spelling)

Telephorus atavinus Scudder, 1891: 587 (unjustified emendation)

Telephorus atavinus Piton & Théobald, 1936: 207

Telephorus atavinus Mitchell 2013 - EDNA The Fossil Insect Database

Cantharis attavina Kirejtshuk & Ponomarenko, 2009-2015

Giebel 1852: 650; Pictet 1854: 334; Giebel 1856a: 102, 400 (in the "Tabellarische"); Giebel, 1856b: 186; Scudder 1876: 81; Goss 1878: 50 (reprint 1878: 328); Scudder 1885: 796; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 587 [under family Lampyridae]; Scudder 1900: 101 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis attavina*]; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis attavina*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Early Miocene: Burdigalian (16-20 Ma).

TYPE LOCALITY. Croatia: Radoboj.

PRESERVATION/LITHOLOGY. Mold / impression/compression (adpression). Lagoonal/restricted shallow subtidal limestone.

COLLECTIONS. Unspecified coll.

COMMENTS AND NOTES. Preserved only abdomen and elytra, that are pales and delicates. Length of fragment: 8.15 mm x 2.50 mm. Similar to the extant *Cantharis livida* Linnaeus, 1758, but smaller. The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331).



Fig. 1 - Cantharidae. Baltic amber. Photo of Artur R.

*¹The original name *attavinus* is present in Latin, consequently the correct name and declination for this species is *attavina* (not *atavina*).

†*Cantharis (Cantharis) brodiei* (C.H.G. von Heyden & L.F.J.D. von Heyden, 1866)

Telephorus Brodiei Heyden & Heyden, 1866: 141, Taf. XXII. Fig. 25. (*separatum*: p. 13, Taf. I. Fig. 25.)

Telephorus Brodieri Piton & Théobald, 1936: 207 (incorrect spelling)

Goss 1878: 55 (reprint 1878: 333); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 587 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis Brodiei*]; Meunier 1915: 220; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis brodiei*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Oligocene (28.4-23.0 Ma). Rott Formation, MP 30.

TYPE LOCALITY. Germany: Rott (Siebengebirge).

PRESERVATION/LITHOLOGY. Adpression/compression. Lacustrine, lithified shale.

COLLECTIONS. Originally in Institut für Paläontologie, Universität Bonn (Krantz coll.).

COMMENTS AND NOTES. Body length: 8.40 mm. Fossilized back, indistinct head. Prothorax narrower than elytra, one conserved antenna to the body side. Elytra more long, wide and dark of the body. Six

visible abdominal segments. The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), but the species is not identifiable given the figure of the descriptors, according Meunier (1915: 220). Specimen known and described by Heyden and Heyden, after Brodie studies.

†*Cantharis (Cantharis) caduca* (C.H.G. von Heyden & L.F.J.D. von Heyden, 1866)

Telephorus caducus Heyden & Heyden, 1866: 141, Taf. XXII. Fig. 20. (*separatum*: p. 13, Taf. I. Fig. 20.)

Goss 1878: 55 (reprint 1878: 333); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 587 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis caduca*]; Meunier 1915: 220; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis caduca*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Oligocene (28.4-23.0 Ma). Rott Formation, MP 30.

TYPE LOCALITY. Germany: Rott (Siebengebirge).

PRESERVATION/LITHOLOGY. Adpression/compression. Lacustrine, lithified shale.

COLLECTIONS. Originally in Institut für Paläontologie, Universität Bonn (Krantz coll.).

COMMENTS AND NOTES. Body length: 9 mm. Head elongated, prothorax narrower than elytra, pale with dark spots: one anterior with two lobes and one small posterior. Short tarsus. Elytra that the leave uncovered



Fig. 2 - Cantharidae. Baltic amber. Photo of Artur R. Michalski



Fig. 3 - Cantharidae. Baltic amber. Photo of Artur R. Michalski

the two last abdominal segments. This species has the shield more smooth of *Cantharis exauctarata*, according to Meunier (1915:220). The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331).

†*Cantharis (Cantharis) carbonaria* (C.H.G. von Heyden & L.F.J.D. von Heyden, 1866)

Telephorus carbonarius Heyden & Heyden, 1866: 140, Taf. XXIII. Fig. 2. (*separatum*: p. 12, Taf. II. Fig. 2.)

Goss 1878: 55 (reprint 1878: 333); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 587 [under family Lampyridae]; Handlirsch 1906-1908: 739 [as *Cantharis carbonaria*]; Meunier 1915: 219, 220-221 (figs. 1-2), Tafel XXVI: Fig. 1; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis carbonaria*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Oligocene (28.4-23.0 Ma). Rott Formation, MP 30.

TYPE LOCALITY. Germany: Rott (Siebengebirge).

PRESERVATION/LITHOLOGY. Adpression/compression. Lacustrine, lithified shale.

COLLECTIONS. Originally in Institut für Paläontologie, Universität Bonn (Krantz coll.).

COMMENTS AND NOTES. Body length: 12 mm, but indicated 20 mm by Meunier (1915: 221). Fossilized back, head ovoidal elongate, prothorax incomplete narrowest of elytra, dark belly, legs all preserved, narrow, long and yellowish. Meunier (1915: 221) redescribes the leg, and particularly the antennomeres, while for the plate provided (Meunier

1915: T. XXVI, Fig. 1), I believe, for the abdomen size and short antennae, that is probably a female. The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), but for the tarsomeres bilobed at sides, the attribution to *Cantharis* is, for me, correct (see Meunier 1915, plate 26 fig. 1).

†*Cantharis (Cantharis) exauctarata* (C.H.G. von Heyden & L.F.J.D. von Heyden, 1866)

Telephorus exauctaratus Heyden & Heyden, 1866: 140-141, Taf. XXIII. Fig. 3. (*separatum*: pp. 12-13, Taf. II. Fig.3.)

Heyden 1870: 265 "Nachträge" (*separatum*: 29 "Nachträge"); Goss 1878: 55 (reprint 1878: 333); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 588 [under family Lampyridae]; Handlirsch 1906-1908: 739 [as *Cantharis exauctarata*]; Meunier 1915: 220; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis exauctarata*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Oligocene (28.4-23.0 Ma). Rott Formation, MP 30.

TYPE LOCALITY. Germany: Rott (Siebengebirge).

PRESERVATION/LITHOLOGY. Adpression/compression. Lacustrine, lithified shale.

COLLECTIONS. Originally in Institut für Paläontologie, Universität Bonn (Krantz coll.), but in coll. Heymann not in Krantz collection, by the same Author (Heyden 1870: 265 "Nachträge"; *separatum*: 29 "Nachträge").

COMMENTS AND NOTES. Body length: 11, 25 mm. Head oblong, slightly more narrow of the prothorax, that is slightly rounded at the sides with vertical black

spotted interrupted in the middle. Parallel elytra and narrow. Thin legs with fourth tarsomer bilobed. The specimen, on the whole, appears black and is preserved also the other plate (negative). The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), but for the prothorax rounded at sides, the attribution to *Cantharis*, is for me correct.

***Cantharis figurata* Mannerheim, 1843**

Cantharis figurata Mann Coope et al., 1961: 390, 410
The Paleobiology Database.

TYPE HORIZON. Late/Upper Pleistocene (about 42.000 years old: 41.500 ± 1200 and 41.900 ± 800). Gottweig (Aurignacian) Interstadial Complex.

TYPE LOCALITY. United Kingdom: Worcestershire, Upton Warren, terrace deposits of the River Salwarpe (Band 3). Latitude $52^{\circ}18'15''N$ - Longitude $2^{\circ}5'40''W$ (National Grid reference SO/935673).

PRESERVATION/LITHOLOGY. Pronotum. Terrestrial, unlithified peat.

COLLECTIONS. Collected in 1950 (The Paleobiology Database).

COMMENTS AND NOTES. One pronotum. The species *Cantharis figurata* is still alive and is an Euro-Asiatic species.

†*Cantharis (Cantharis) fragilis* (Heer, 1847)

Telephorus fragilis Heer, 1847: 148-149, 225, Taf. IV. Fig. 14., Fig. 14. bc.
Giebel 1852: 650; Pictet 1854: 334; Giebel 1856a: 102, 400 (in the "Tabellarische"); Giebel 1856b: 185; Scudder 1876: 81; Goss 1878: 48 (reprint 1878: 326); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 588 [under family Lampyridae]; Scudder 1900: 101 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis fragilis*]; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; ETH Zürich: [20]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis fragilis*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database

TYPE HORIZON. Upper/Middle Miocene: Upper Tortonian/Lower Sarmatian (10-12.7 Ma), Upper Freshwater-Molasse Formation.

TYPE LOCALITY. Germany: Baden-Württemberg, Oeningen (= Öhningen).

PRESERVATION/LITHOLOGY. Mold impression/compression (adpression). Lacustrine, lithified limestone.

COLLECTIONS. Carlsruhe collection, in the original description. Department of Earth Sciences, Inv. No. Pl. I. 292, ID 2606 and 2989 (see ETH Zürich).

COMMENTS AND NOTES. Eutra black. Body length: 7. 90 mm without the head that is not preserved; elytra around 5 mm (body length: 7.40 mm, and elytra: 4.80 mm in The Paleobiology Database) and similar to the extant *C. flavilabris* Fallén, 1807 (*Telephorus fulvicollis* in the description).

The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), but, for me, may remain in *Cantharis* because (although in the figure 14. c., does not appear, and seems similar to *Rhagonycha germari* and *Rh. tertaria*), Heer says in the original description that the third tarsomer is expanded to the outside, so it may be also, for me, bilobed as in the true *Cantharis*.

†*Cantharis (Cantharis) macilenta* (Heer, 1865)

Telephorus macilentus Heer, 1865: Fig. 256., Fig. 256. bcd. (p. 376), 377
Heer 1872: Fig. 256., Fig. 256. bcd. (p. 461), 462; Heer 1876: Fig. 256., Fig. 256. bcd (p. 32), 33; Goss 1878: 48 (reprint 1878: 326); Heer 1883: Fig. 298., Fig. 298. bcd. (p. 402), 403; Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 588 [under family Lampyridae]; Scudder 1900: 101 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis macilenta*]; Handlirsch 1920-1921: 232; ETH




Fig. 4 - *Malthodes* sp. Baltic amber. Photo of Artur R. Michalski

Zürich: [20]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis macilenta*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Upper/Middle Miocene: Upper Tortonian/Lower Sarmatian (10-12.7 Ma), Upper Freshwater-Molasse Formation.

TYPE LOCALITY. Germany: Baden-Württemberg, Oeningen (= Öhningen).

PRESERVATION/LITHOLOGY. Mold impression/compression (adpression). Lacustrine, lithified limestone.

COLLECTIONS. Collection of Zürich University (see The Paleobiology Database). Department of Earth Sciences, Inv. No. Pl. I. 293 and 293b, ID 2607 and 2608 (see ETH Zürich).

COMMENTS AND NOTES. From the original work (and subsequent reprints), we have only the drawing of the species and legs (the posterior legs are swollen), but nothing else details of size, prothorax shape or antennae (only the tibiae are light colored). The bilobed third article of tarsomeres is fairly typical of the genus *Cantharis*.

†*Cantharis aff. nigricans* Burmeister, 1832

Cantharis nigricans Fabr. sehr ähnlich, Burmeister, 1832: 635
Keferstein 1834: 329; Vollmar 1835: 61; Burmeister 1836: 577;
Giebel 1856a: 102; Scudder 1891: 483 [compared with the family Meloidae]; Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Collection in Berlin.

COMMENTS AND NOTES. One specimen.

***Cantharis cf. paludosa* Fallén, 1807**

Cantharis cf. paludosa Coope et al., 1961: 390

The Paleobiology Database.

TYPE HORIZON. Late/Upper Pleistocene (about 42.000 years old: 41.500 ± 1200 and 41.900 ± 800). Gottweig (Aurignacian) Interstadial Complex.

TYPE LOCALITY. United Kingdom: Worcestershire, Upton Warren, terrace deposits of the River Salwarpe (Band 3). Latitude 52°18'15"N - Longitude 2°5'40"W (National Grid reference SO/935673).

PRESERVATION/LITHOLOGY. Head. Terrestrial, unlithified peat.

COLLECTIONS. Collected in 1950 (The Paleobiology Database).

COMMENTS AND NOTES. One head. The species *Cantharis paludosa* is still alive and is an European-Siberian species.

***Cantharis rufa* Linnaeus, 1758**

C. rufa L. (*liturata* Fall.) Coope et al., 1961: 390, 402 (Explanation of plate 21), plate 21 figure 62. [pronotum as *Cantharis liturata*]

The Paleobiology Database.

TYPE HORIZON. Late/Upper Pleistocene (about 42.000 years old: 41.500 ± 1200 and 41.900 ± 800). Gottweig (Aurignacian) Interstadial Complex.

TYPE LOCALITY. United Kingdom: Worcestershire, Upton Warren, terrace deposits of the River Salwarpe (Band 3). Latitude 52°18'15"N - Longitude 2°5'40"W (National Grid reference SO/935673).

PRESERVATION/LITHOLOGY. Various head and pronotum. Terrestrial, unlithified peat.

COLLECTIONS. Collected in 1950 (The Paleobiology Database).

COMMENTS AND NOTES. 26 head, 21 pronotum, and? left elytron. The species *Cantharis rufa* is still alive and is an Euro-Asiatic species, introduced also in North America.

†*Cantharis (Cantharis) sucinonigra* Kuśka, 1992

Cantharis sucinonigra Kuśka, 1992: 107-109, (figs. 1-3 p. 108)

Cantharis succinonigra Alekseev, 2013: 8 (incorrect spelling)

Cantharis succinonigra Kirejtshuk & Ponomarenko, 2009-2015 (incorrect spelling)

Kuśka 1994: 6; Kuśka 1996b: 13; Kosmowska-Ceranowicz et al. 2001: 61; Pawłowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 287, 290; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of the Earth PAS in Warsaw (Museum Ziemi collection), No. 18094.

COMMENTS AND NOTES. Male, with elytra pitchy black and pronotum red. Body length: 6.0 mm, width of pronotum 1.5 mm. Similar to the extant *C. nigra* De Geer.

†*Cantharis* sp. (Hope, 1836)

Necydalis (sic!) Hope, 1836: 143

Berendt 1845: 56 [as ?*Necydalis*]; Giebel 1852: 656 [as *Necydalis*]; Scudder 1885: 793 [as *Necydalis*]; Scudder 1887: 794 [as *Necydalis*]; Handlirsch 1906-1908: 787 [as

Fig. 5 - Myamalycoerus vitalii Fanti & Ellenberger, 2016 - courtesy of Sieghard Ellenberger





Fig. 6 -*Tytthonyx geiseri* Photo courtesy of Poinar

Necydalis ?, in the family Cerambycidae]; Spahr 1981b: 15; Poinar 1992: 138; Douglas & Stockey 1996: 1154; Vitali 2009: 236 [as probably a “*Cantharis*”].

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. ? Berendt collection.

COMMENTS AND NOTES. Probably a *Cantharis*, in fact, according to Vitali (2009: 236) no *Necydalis* (Coleoptera, Cerambycidae) is known for the Baltic amber.

†*Cantharis* sp. (Hope, 1836)

Telephorus Hope, 1836: 143

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Hope authority.

†*Cantharis* sp. Berendt, 1845

Cantharis Berendt, 1845: 47, 56

Hope 1836: 143 (Berendt authority, as *Cantharis* and *Telephorus*); Pictet 1846: 103 [as *Telephorus*]; Pictet 1854: 334; Giebel 1856a: 102; Giebel, 1856b: 181 [as *Telephorus*]; Fischer 1939: 97 [as *Cantharis*?], (p. 96 as *Cantharis rubiniformis minima*, but it is not Cantharidae); Larsson 1978: 139; Spahr 1981b: 4, 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Berlin Museum (coll. Berendt).

COMMENTS AND NOTES. 9 specimens. Almost one of these was firstly mentioned by Hope (1836: 143).

†*Cantharis* sp. (Berendt, 1845)

Molorchus Berendt, 1845: 56

Giebel 1852: 656 [as *Molorchus*]; Giebel 1856: 128, 403 (in the “Tabellarische”) [as *Molorchus*]; Scudder 1885: 793 [as *Molorchus*]; Scudder 1887: 794 [as *Molorchus*]; Zang 1905: 233 [as *Cantharis*]; Handlirsch 1906-1908: 739 [as ? *Cantharis* sp.]; Handlirsch 1920-1921: 232; Larsson 1978: 139; Spahr 1981b: 15; Poinar 1992: 138; Douglas & Stockey 1996: 1154; Kazantsev 2013: 283; Vitali 2009: 237; Kirejtshuk & Ponomarenko 2009-2015

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber.

Baltic Amber.

COLLECTIONS. Berlin Museum (coll. Berendt).

COMMENTS AND NOTES. Specimen examined by Zang (1905: 233) resulting to be a *Cantharis*-species with damaged elytra (Zang 1905: 233; Vitali 2009: 237).

†*Cantharis* sp. Giebel, 1852

Cantharis Giebel, 1852: 655

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. One specimen.

†*Cantharis* sp. (Giebel, 1856)

Telephorus Giebel, 1856a: 400 (in the “Tabellarische”)

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. 9 specimens.

†*Cantharis* sp. Motschulsky, 1857

Cantharis Motschulsky, 1857: 28 [as *Cantharis*], 30 [as *Oripa* (*Cantharis*)]

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Coll. Menge.

COMMENTS AND NOTES. One or more specimens, and one of this as *Oripa* and longer than the extant *Oripa dispar* = now *Cantharis livida* Linnaeus (Motschulsky, 1857: 30), but it is uncertain whether in coll. Menge or Berendt.

†*Cantharis* sp. Helm, 1896

Cantharis Helm, 1896: 228

Handlirsch 1906-1908: 739; Handlirsch 1920-1921: 232; Larsson 1978: 139; Spahr 1981b: 15; Kazantsev 2013: 283; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Cantharis* sp. Klebs, 1910



Figs. 7-8: *Cacomorphocerus* sp. Baltic amber - courtesy of Anders L. Damgaard.
Fig. 9: Cantharidae. Baltic amber - courtesy of Anders L. Damgaard.





Fig. 10: *Ornatomalthinus elviraе* Poinar & Fanti, 2016. Photo of G. Poinar

Fig. 11: *Rhagonycha sucinobaltica* Poinar & Fanti, 2016. Photo of G. Poinar

Cantharis Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 15; Hieke & Pietrzeniuk 1984: 303, 321; Poinar & Fanti 2016: 1.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection, collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Twelve specimens, another one near to *Cantharis*, and one between *Cantharis* and *Rhagonycha*.

†***Cantharis* sp.** (Piton & Théobald, 1936)

Telephorus sp. Piton & Théobald, 1936: 206-207, 211, 212, 213 Fig. 10.

The Paleobiology Database.

TYPE HORIZON. Late/Upper Oligocene (28.4-23.3 Ma) or Early Miocene (about 22.0 Ma).

TYPE LOCALITY. France: near of Clermont-Ferrand, right bank of Allier, North-East of Puy-de-Mur, deposit of Puy-Saint-Jean.

PRESERVATION/LITHOLOGY. Elytra. Crate lake limestone (volcanic tuff with intercalations of the calcareous marl).

COLLECTIONS. M. A. Rudel collection.

COMMENTS AND NOTES. Two specimens. Length: 5 mm. Elytron light yellow, white spot near scutellum, blackish tint to the top. Similar to the extant *Cantharis livida* var. *bicolorata* Ragusa.

†***Cantharis* sp.** Bachofen-Echt, 1949

Cantharis Bachofen-Echt, 1949: 109

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†***Cantharis* sp.** Hieke & Pietrzeniuk, 1984

Cantharis Hieke & Pietrzeniuk, 1984: 303

Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museums für Naturkunde, Berlin.

COMMENTS AND NOTES. Hieke & Pietrzeniuk (1984: 303, 321) quote the genera: *Cantharis*, *Rhagonycha*, *Malthinus* and *Malthodes*, studying about 110 specimens of various Museums and collections: Berlin: Künnow collection - 25 specimens, Berendt collection - 10 specimens, Kühl collection - 7 specimens, Simon collection - 3 specimens, Isenheim collection - 3 specimens, "Bitterfelder Bernstein" - 1 specimen, Mineralogischen Museum - 1 specimen; Klebs collection - 47 specimens; Copenhagen collection - ? few.

†***Cantharis* sp.** Schawaller, 1986

Cantharis nahe / *Cantharidae* Schawaller, 1986: 3, 5 (Abb. 9 [specimen No. 61205])

The Paleobiology Database.

TYPE HORIZON. Early-Middle Miocene: Burdigalian/Karpatian (16.0-13.7 Ma).

TYPE LOCALITY. Germany: Baden-Württemberg, Randecker Maar.

PRESERVATION/LITHOLOGY. Mold/impression. Crater lake, carbonaceous lime mudstone.

COLLECTIONS. Staatlichen Museum für Naturkunde Stuttgart, Nr. 61201 to 61205.

COMMENTS AND NOTES. Five specimens. One (specimen No. 61205) with pronotum probably red (elytra and legs black?) and about 9 mm.

Subgenus *Cyrtomoptila* Motschulsky, 1860

†***Cantharis* (*Cyrtomoptila*) *sucinokotejai*** (Kuška, 1996)

Absidiella sucinokotejai Kuška, 1996b: 14-15, 18 (fig. 7)

Absidiella sucinokotejai Pielińska, 2006: 9, english part: 34 (incorrect spelling)

Albsidiella succinokotejai Alekseev, 2013: 8 (incorrect spelling)

Albsidiella ??? *succinokotejai* Kirejtshuk & Ponomarenko, 2009-2015 (incorrect spelling)

Kuška 1994: 3, 6 (only as *Absidiella* n. sp., No. 10424, female, Museum of the Earth collection; and a male in poor condition indicated as *Cantharis* sp. (p. 3) or as *Absidiella* (p. 6); Kuška, 1996a: 19 (species unnamed); Kosmowska-Ceranowicz et al. 2001: 61 [as *Absidiella sucinokotejai*]; Pawłowski & Mazur 2012: 44 [as *Absidiella sucinokotejai*]; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database [as *Absidiella sucinokotejai*]; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.
 PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.
 COLLECTIONS. Museum of the Earth PAS in Warsaw (Museum Ziemi collection), No. 10424.
 COMMENTS AND NOTES. Female, brown, thorax and femora black-brown, tibiae and tarsi yellow-brown. Body length: 6.0 mm, elytra length 4.25 mm and width 1.5 mm, elytra index 2.8 mm. Related to the extant *Podistra schoenherri* Dejean. Transferred to *Cantharis* (*Cyrtomoptila*) by Kazantsev (2013: 290). A male of the same genus *Absidiella* No. 10424 is present in Museum of the Earth in Warsaw but is in a poor condition (Kuśka, 1994: 6).

Genus †*Electronycha* Kazantsev, 2013: 284-285, 290

†*Electronycha prussica* Kazantsev, 2013
Electronycha prussica Kazantsev, 2013: figs. 1-3 (p. 284), 285, 290
 Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; The Paleobiology Database.
 TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).
 TYPE LOCALITY. Russian Federation: Baltic Sea coast.
 PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.
 COLLECTIONS. Senckenberg Deutsches Entomologisches Institut (Müncheberg - Germany), No. 1112-3.
 COMMENTS AND NOTES. Male, dark brown. Body length: 7.4 mm, width 2.0 mm. Genus characterized to 15-segmented antennae: 1-5 and 10-15 filiforms, and 6-9 distally swollen. Syninclusions: various.

Genus †*Lithocantharis* Lin, 1997: 186, 188, 189, 193, 194, 199 (caption)

†*Lithocantharis lunglokshuiensis* Lin, 1997
Lithocantharis lunglokshuiensis Lin, 1997: 186, 188, 193 (fig. 5), 199 [caption of the plate 3], pl. 3 fig. 1
 The Paleobiology Database.
 TYPE HORIZON. Paleocene (66-56 Ma). Ping Chau Formation of China.
 TYPE LOCALITY. Hong Kong (China), Peng Chau Island, (also known as Tung Ping Chau Island), Lung Lok Shui (LLT).
 PRESERVATION/LITHOLOGY. Elytron (compression). Lithified, terrestrial siltstone.
 COLLECTIONS. Nanjing Institute of Geology and Palaeontology (NIGP), No. 128290 (LLT-1-4).

COMMENTS AND NOTES. It is preserved only an elytron, length: 5.0 mm x 2.0 mm. Similar to *Cantharis* but with wings darker and narrower. For the presence of various striae on elytra, membership in the Cantharidae family is, for me, uncertain.

Genus *Lycocerus* Gorham, 1889

†*Lycocerus guttula* (J. Zhang, 1989) [*n. comb.*]
Cantharis guttula J. Zhang, 1989: 119 (illustration 103)-120, 387, 445 (caption of Table 26), Table 26 Fig. 4.
 Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Middle Miocene (16.0-11.6 Ma). Shanwang Formation.

TYPE LOCALITY. China: Shandong, Shanwang, Linqu County, collection 3.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large mudstone.

COLLECTIONS. Shandong Provincial Museum (LPM), No. 830063.

COMMENTS AND NOTES. Head and wings brown dark, prothorax stained. Body length: 8.80 mm x 2.90 mm. Zhang says that is similar to *Cantharis oedemeroides* Kiesenwetter = *Lycocerus oedemeroides* (Okushima 2005: 283). For this similarity, for the general appearance, and particularly for slender body and for the prothorax sub-quadrata that is typical on *Lycocerus oedemeroides* species-group (Okushima 2005: 273; Hsiao & Okushima 2015: 109), I tentatively transferred this species to genus *Lycocerus* Gorham.

Genus †*Myamalycopterus* Fanti & Ellenberger, 2016: 166-167

†*Myamalycopterus vitalii* Fanti & Ellenberger, 2016
Myamalycopterus vitalii Fanti & Ellenberger, 2016: 167-169 (figs. 1-3)
 The Paleobiology Database.

TYPE HORIZON. Early Cretaceous: Upper Albian (97-110 Ma).

TYPE LOCALITY. Myanmar: Kachin State, Myitkyina District, Tanai Township, Hukawng Valley, Aung Bar Maw mine.

PRESERVATION/LITHOLOGY. Inclusion in amber. Burmese Amber.

COLLECTIONS. SMNS Naturkunde Museum Stuttgart (Germany), SMNS BU-230.

COMMENTS AND NOTES. Similar to *Ornatomalthinus elvira* Poinar & Fanti, but differs by the long elytra and relief points which are less raised and more numerous.

Genus †*Ornatomalthinus* Poinar & Fanti, 2016: 2

†*Ornatomalthinus elviraе* Poinar & Fanti, 2016

Ornatomalthinus elviraе Poinar & Fanti, 2016: 2-3 (figs. 1-3), 5-6
 Poinar et al. 2007: 1663-1668 (figs. 1abc, 2abc, 3); Binder 2008: 47 (Abb. 1); Oktar / [Harun Yahya] 2006-2008: 612-613 (figs.); Boucot & Poinar 2010: 196 (Figure 269); Ross et al. 2010: 217; Rasnitsyn et al. 2016: Appendix A, Appendix B p. 7; Hsiao et al. 2016: 120, 122; Fanti & Ellenberger 2016: 166, 168, 169; The Paleobiology Database.

TYPE HORIZON. Early Cretaceous: Upper Albian (97-110 Ma).

TYPE LOCALITY. Myanmar: Kachin State, Southwest of Maingkhwan, Hukawng Valley, mine near Noije Bum, designated as "Noije Bum 2001 Summit Site" (26°20'N, 96°36'E).

PRESERVATION/LITHOLOGY. Inclusion in amber. Burmese Amber.

COLLECTIONS. George Poinar amber collection at Oregon State University, Corvallis (USA), No. B-C-28. Another similar specimen (Poinar et al. 2007) actually is in the Deniz Eren amber collection, Istanbul (see down).

COMMENTS AND NOTES. Female with short elytra with evident striation equipped with relief points. Body length: 3.5 mm. Genus characterized for possessing features similar to lycid beetles. Another specimen in Burmese amber from the "Noije Bum 2001 Summit Site" and deposited in the amber collection of Ron Buckley No. ABS66; long about 4.5 mm and with extruded cuticular vesicle glands, is illustrated by Poinar et al. (2007: 1663-1668) and after reillustrated in Binder (2008: Abb. 1), Oktar / [Harun Yahya] (2006-2008: 612-613 [same specimen?]), and Boucot & Poinar (2010: Fig. 269, specimen in the Deniz Eren amber collection, Istanbul - Turkey). It has similar characteristics but without further studies it is not possible to say if is related to *Ornatomalthinus elviraе* (Poinar & Fanti, 2016: 5-6).

Genus *Podistra* Motschulsky, 1839**Subgenus *Absidia* Mulsant, 1862****†*Podistra (Absidia)* sp. (Klebs, 1910)**

Absidia Klebs, 1910: 237

Handlirsch 1920-1921: 232 [as *Absidia*]; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 15; Hieke & Pietrzeniuk 1984: 303, 321; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection,

collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Two specimens.

†*Podistra (Absidia)* sp. (Bachofen-Echt, 1949)

Absidia Bachofen-Echt, 1949: 109

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

Genus *Rhagonycha* Eschscholtz, 1830**Subgenus (*Rhagonycha*) Eschscholtz, 1830****†*Rhagonycha (Rhagonycha) germari* (Heer, 1847) [n. comb.]**

Telephorus Germari Heer, 1847: 143-145, 225, Taf. IV. Fig. 10., Fig. 10. bcde.

Telephorus Germarii Heer, 1847: 225 ([caption] Fig. 10.); Taf. IV. ([caption] (Fig. 10.) (incorrect original spelling. ICZN 1999 Art. 19.3)

Telephorus Germarii Scudder, 1876: 81 (incorrect subsequent spelling. ICZN 1999 Art. 33.4)

Giebel 1852: 650; Pictet 1854: 334; Giebel 1856a: 101, 400 (in the "Tabellarische"); Giebel, 1856b: 185; Scudder 1876: 81; Goss 1878: 48 (reprint 1878: 326); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 588 [under family Lampyridae]; Scudder 1900: 101 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis Germari*]; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207; Carpenter 1992: 331 [as *Cantharis*]; ETH Zürich: [20]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis germari*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Upper/Middle Miocene: Upper Tortonian/Lower Sarmatian (10-12.7 Ma), Upper Freshwater-Molasse Formation.

TYPE LOCALITY. Germany: Baden-Württemberg, Oeningen (= Öhningen).

PRESERVATION/LITHOLOGY. Mold impression/compression (adpression). Lacustrine, lithified limestone.

COLLECTIONS. Two specimens in Carlsruhe collection, in the original description. Department of Earth Sciences, Inv. No. Pl. I. 290, ID 2604 (see ETH Zürich).

COMMENTS AND NOTES. Elytra black. Body

length: 9.0 mm ("4 Lin.") x 2.80 mm, but of the holotype is preserved only elytra and abdomen (Fig. 10.). Length sometimes indicated as 8.50 x 2.60 mm (The Paleobiology Database). Other specimen of the original description is more well preserved (Fig. 10. c.). Similar to the extant *C. rustica* and *C. fusca*, especially *C. rustica* by Giebel (1856: 101), but the assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), in fact, particularly, for the original picture (Taf. IV. Fig. 10. d.) and the description of the tarsus with third tarsomer non bilobed, I tentatively transferred this species to the genus *Rhagonycha* Eschscholtz, 1830.

†*Rhagonycha (Rhagonycha) hesperus* (Wickham, 1914) [n. comb.]

Telephorus hesperus Wickham, 1914: 445, Plate 4. Fig. 3. (and explanation) [under family Lampyridae]

International Catalogue of Scientific Literature 1916b: 148; International Catalogue of Scientific Literature 1919: 108; Meunier 1920a: 156; Wickham 1920: 353 [as *Cantharis hesperus*]; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Cantharis*]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis hespera*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database [as *Cantharis hesperus*].

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale, volcaniclastic.

COLLECTIONS. Holotype: Museum of Comparative Zoology - Harvard University (MCZ), No. 2,496 (No. 9,376 S. H. Scudder coll.). Other specimens MCZ, No. 2,497 to 2,501 (No. 2,243, 5,065, 5,515, 6,048, 12,769 S. H. Scudder coll.).

COMMENTS AND NOTES. Yellowish, with sides and sutural region of the elytra darker. Body length: 4.50 mm to apex of elytra, other specimens little over 5 mm. Described from six specimens. In the original description is indicated as similar to the recent *Telephorus scitulus* (= *Rhagonycha scitula* (Say), and for this and for the little length of the all specimens, I tentatively transferred to genus *Rhagonycha*, although the rounded prothorax leaves doubt. The assignment to genus *Cantharis* was doubt also for Carpenter (1992: 331). The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

†*Rhagonycha (Rhagonycha) kryshtofovichi* (Yablokov-Khnzorian, 1960)

Malchinus kryshtofovichi Yablokov-Khnzorian, 1960: fig. 3abvg (p. 94), 95

Larsson 1978: 140; Spahr 1981b: 15; Poinar 1992: 137; Kuška

1994: 6; Douglas & Stockey 1996: 1154; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1, 4, 6; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Lithuania: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Paleontological Institute, Russian Academy of Sciences (PIN) in Moscow, No. 364/108.

COMMENTS AND NOTES. Male, reddish brown, pronotum slightly narrowed anteriorly. Body length: 7 mm. Transferred to *Rhagonycha* by Kazantsev (2013: 290).

†*Rhagonycha (Rhagonycha) micans* Piton, 1939

Rhagonycha micans Piton, 1939: 102, Planche I fig. 6 (p. [104]), 105

Poinar & Fanti 2016: 6; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Pliocene: Piacenzian (3.6-2.6 Ma).

TYPE LOCALITY. France: Puy-de-Dôme, Lac Chambon.

PRESERVATION/LITHOLOGY. Elytra and pronotum (impression/compression fossil). Crater lake, diatomite and volcaniclastic.

COLLECTIONS. L. Piton collection, No. 1023 (holotype). Probably redeposited at MNHN (Paris). Piton legit in 1938.

COMMENTS AND NOTES. Two elytra and fragments of the pronotum. Elytra red very clear with reflections iridescent, and with fine striae punctate. Near the extant *Rhagonycha testacea* (Linnaeus, 1758), but with end of the elytra more narrowed.

†*Rhagonycha (Rhagonycha) sucinobaltica* Poinar & Fanti, 2016

Rhagonycha sucinobaltica Poinar & Fanti, 2016: 3-4 (figs. 4-6), 6

Hsiao et al. 2016: 120; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Kaliningrad region, Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. George Poinar amber collection at Oregon State University, Corvallis (USA), No. C-202.

COMMENTS AND NOTES. Male, blackish to dark

brown with pronotum slightly darker than elytra. Body length: 3.5 mm. Similar to the extant *Rhagonycha atra* group.

†*Rhagonycha (Rhagonycha) tertaria* (Heer, 1847)
[n. comb.]

Telephorus tertarius Heer, 1847: 145-148, 225, Taf. IV. Fig. 11, 11bc., 12, 12bc., 13

= *Telephorus tertarius oeningensis* Heer, 1847: 145-146, 225, Taf. IV. Fig. 11., Fig. 11. bc. [n. syn.]

= *Telephorus tertarius Radobojanus* Heer, 1847: 147-148, 225, Taf. IV. Fig. 12., Fig. 12. bc. [n. syn.]

Giebel 1852: 650 [as *Telephorus tertarius*]; Pictet 1854: 334 [as *Telephorus tertarius* (*sic!*)]; Giebel 1856a: 101-102, 400 (in the “Tabellarische”) [as *Telephorus tertarius*]; Giebel, 1856b: 185, 186 [as *Telephorus tertarius*]; Scudder 1876: 81; Goss 1878: 48, 50 (reprint 1878: 326, 328); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 588 [as *Telephorus tertarius*, *Telephorus tertarius oeningensis* and *Telephorus tertarius radobojanus* (*Telephorus oeningensis* and *Telephorus radobojanus*); under family Lampyridae]; Scudder 1900: 101 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Cantharis tertaria radobojana* and *Cantharis tertaria oeningensis*]; Handlirsch 1920-1921: 232; Piton & Théobald 1936: 207 [as *Telephorus radobojanus* and *Telephorus tertarius*]; Carpenter 1992: 331 [as *Cantharis*]; ETH Zürich: [20]; Kirejtshuk & Ponomarenko 2009-2015 [as *Cantharis radobojana*, *Cantharis tertaria* and *Cantharis tertaria oeningensis*]; Mitchell 2013 - EDNA The Fossil Insect Database [as *Telephorus tertarius* (*sic!*) *radobojanus* and *Telephorus tertarius oeningensis*]; The Paleobiology Database [as *Telephorus tertarius*].

TYPE HORIZON. Upper/Middle Miocene: Upper Tortonian/Lower Sarmatian (10-12.7 Ma), Upper Freshwater-Molasse Formation (for the Oeningen specimen). [Early Miocene: Burdigalian (16-20 Ma) for the Radoboj specimen].

TYPE LOCALITY. Germany: Baden-Württemberg, Oeningen (= Öhningen). [ICZN 1999 Art. 76.2.]. The other specimen comes from Croatia: Radoboj.

PRESERVATION/LITHOLOGY. Mold impression/compression (adpression). Lacustrine, lithified limestone.

COLLECTIONS. Unspecified coll. Department of Earth Sciences, Inv. No. Pl. I. 291, ID 2605 (see ETH Zürich) for Oeningen specimen.

COMMENTS AND NOTES. Elytra testaceous with the black apex, and similar (also for the original description) to the extant *Telephorus melanura* F. = *Rhagonycha fulva* (Scopoli, 1763); with thinner antennae by Giebel (1856: 101). Body length: around 7.25 mm. (7.15 mm x 2.10 mm by The Paleobiology Database).

The assignment to genus *Cantharis* is doubt for Carpenter (1992: 331), in fact, particularly, for the original pictures (Taf. IV. Figs. 11. c., 12. bc.) and the descriptions of the tarsus with third tarsomer

non bilobed, and also for the narrowed pronotum, I tentatively transferred this species to the genus *Rhagonycha* Eschscholtz, 1830.

Heer briefly described the species *Telephorus tertarius*, but having two specimens of two different locations and especially not kept fully, described these exemplars with two different names at subspecies level (*T. t. oeningensis* and *T. t. radobojanus*), mistakenly leaving out the nominotypical subspecies. But in the original description of the two subspecies, there are not substantial differences, in fact, have the exact same coloration, same shape, same legs (tarsus) and almost the same size. Actually also geographically the different subspecies are not justified, but however, the fossil faunas of the two places are not presently considered coeval, which leaves a minimum doubt about synonymies here presented. Moreover, the subspecies have the same type of *T. tertarius*, therefore are to be considered objective synonyms (ICZN 1999 Art. 61.3.4).

The species is therefore composed of two syntypes and it is necessary to designate the lectotype. Seen that the Fig. 11. (Taf. IV. Fig. 11. Specimen of Oeningen) is the first to appear in the original description and above the specimen is slightly more complete fossilized (are missing part of the head and prothorax included some legs), is here considered as lectotype, and Fig. 12. (Taf. IV. Fig. 12. Specimen of Radoboj) as paralectotype (ICZN 1999 Art. 74.1.3., Art. 74.4., Recommendation 74F.).

†*Rhagonycha* sp. Helm, 1896

Rhagonycha Helm, 1896: 228

Handlirsch 1906-1908: 739; Handlirsch 1920-1921: 232; Larsson 1978: 139; Spahr 1981b: 16; Kazantsev 2013: 283.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Rhagonycha* sp. Klebs, 1910

Rhagonycha Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 16; Hieke & Pietreniuk 1984: 303, 321; Poinar & Fanti 2016: 1, 6; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection,

collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Fifteen specimens, and five near to *Rhagonycha*, and one between *Cantharis* and *Rhagonycha*.

†*Rhagonycha* sp. Bachofen-Echt, 1949

Rhagonycha Bachofen-Echt, 1949: 109

Spahr 1981b: 16.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Rhagonycha* sp. Hieke & Pietrzeniuk, 1984

Rhagonycha Hieke & Pietrzeniuk, 1984: 303

Poinar & Fanti 2016: 1, 6; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museums für Naturkunde, Berlin.

COMMENTS AND NOTES. Hieke & Pietrzeniuk (1984: 303, 321) quote the genera: *Cantharis*, *Rhagonycha*, *Malthinus* and *Malthodes*, studying about 110 specimens of various Museums and collections: Berlin: Künow collection - 25 specimens, Berendt collection - 10 specimens, Kühl collection - 7 specimens, Simon collection - 3 specimens, Isenheim collection - 3 specimens, "Bitterfelder Bernstein" - 1 specimen, Mineralogischen Museum - 1 specimen; Klebs collection - 47 specimens; Copenhagen collection - ? few.

Genus †*Sucinocantharis* Kuška & Kania, 2010:

52

†*Sucinocantharis baltica* Kuška & Kania, 2010

Sucinocantharis baltica Kuška & Kania, 2010: 52-53 (figs. 7-9)

Pawlowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kania 2015: 12; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4

Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Deutsches Entomologisches Institut (DEI) in Germany, No. 1441 - 1 (ex coll. C. and H. W. Hoffeins, Hamburg).

COMMENTS AND NOTES. Male. Black. Body length: 8.0 mm. Genus characterized to 16-segmented antennae, with antennomeres 3-10 dilated. Syninclusions: Diptera Bibionidae.

Genus †*Sucinorhagonycha* Kuška, 1996: 13

†*Sucinorhagonycha kulickae* Kuška, 1996

Sucinorhagonycha kulickae Kuška, 1996b: 13-14 (figs. 1-2), 17 (figs. 3-5)-18 (fig. 6)

Kuška 1994: 3, 6 (only as *Rhagonycha* n. sp., No. 22345, male, Museum of the Earth collection); Kuška, 1996a: 19 (species unnamed); Kubisz 2000: 226, 227-229 (fig. 1 p. 228); Kosmowska-Ceranowicz et al. 2001: 61; Kuška & Kania 2010: 49; Pawłowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Holotype: Museum of the Earth PAS in Warsaw (Museum Ziemi collection), No. 22345. Female described to Kubisz is preserved in Museum of Natural History of the Institute of Systematics and Evolution of Animals (ISEA) in Kraków, No. 1/31.

COMMENTS AND NOTES. Male, black-brown with pronotum and elytra with a distinct sheen. Body length: 4.5 mm (elytra 3.5 mm). Genus characterized to 12 filiform antennomeres. Female (Kubisz 2000: 227-229) with black elytra and body length 5 mm.

Genus *Themus* Motschulsky, 1858

= †*Curticantharis* J. Zhang, 1989: 120, 422 [*n. syn.*] *²

*²The genus *Curticantharis* according to Zhang, is characterized for the head short, pronotum subquadrate/rectangular with straight edges and slightly rounded corners, short filiform antennae 11-segmented, small scutellum, short legs, large size, and particularly the elytra which do not cover the abdomen. But these characters are in communion with other genera and in detail, in area in question, with *Themus* (especially

pronotum sub-quadrata/transverse, relative short antennae and large size), which is a genus widespread in China and Asia. The character believed important, of the short elytra, may be due, in the Zhang species, for an abnormal distension during fossilisation, and in more the females of many genera have the wide body and stretched beyond of the wing-covers and/or physogastric (common in the females of *Themus*). Finally the three species of *Curticantharis* have a different look: broad and squat or slender elongated, compatible with the diversity of habitus of *Themus*. All *Curticantharis* species of Zhang, I tentatively placed to nominotypical subgenus *Themus* for the lateral margins of pronotum straight, parallel-sided (see: Švihla 2008; Yang & Yang 2010).

Subgenus *Haplothemus* Wittmer, 1973

†*Themus (Haplothemus) pristinus* Kazantsev, 2013

Themus pristinus Kazantsev, 2013: 286-288 (figs. 7-9 p. 287), 290

Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Senckenberg Deutsches Entomologisches Institut (Müncheberg - Germany), No. 1593-1.

COMMENTS AND NOTES. Female, dark brown. Body length: 13.2 mm, width 3.2 mm. Species characterized for large size and long cheeks. Syninclusions: one Diptera.

Subgenus *Themus* Motschulsky, 1858

†*Themus (Themus) capacis* (J. Zhang, 1989) [n. comb.]

Curticantharis capacis J. Zhang, 1989: 120 (and key)-121 (illustration 104 A.B.C.D.), 122, 387, 422, 445 (caption of Table 26), Table 26 Fig. 5., (8 [as Cantharidae])

Zhang et al 1994: 5, 6, 9, 265 [as Cantharidae]; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Middle Miocene (16.0-11.6 Ma). Shanwang Formation.

TYPE LOCALITY. China: Shandong, Shanwang, Linqu County, collection 3.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large mudstone.

COLLECTIONS. Shandong Provincial Museum (LPM), No. 820289.

COMMENTS AND NOTES. Body length: 15.4 mm x 5.4 mm. Species smallest, black, with more broad wings than *Themus thermophilus*.

†*Themus (Themus) thermophilus* (J. Zhang, 1989)

[n. comb.]^{*3}

Curticantharis thermophila J. Zhang, 1989: 120 in key, 121 (illustration 105)-122, 387, 445 (caption of Table 27), Table 27 Fig. 3., (8 [as Cantharidae])

Zhang et al. 1994: 5, 6, 9, 265 [as Cantharidae], 90-91; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Middle Miocene (16.0-11.6 Ma). Shanwang Formation.

TYPE LOCALITY. China: Shandong, Shanwang, Linqu County, collection 5.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large mudstone.

COLLECTIONS. Shandong Provincial Museum (LPM), No. 750097 and 750096.

COMMENTS AND NOTES. Two specimens. Body length: 19.5 mm x 6.8 mm. Body more wide and reddish-brown with narrow wing than *Themus capacis*.

^{*3}The female name *thermophila* was granted by me from the masculine genus *Themus* in *thermophilus*.

†*Themus (Themus) trapezialis* (J. Zhang, Sun & X. Zhang, 1994) [n. comb.]

Curticantharis trapezialis J. Zhang, Sun & X. Zhang, 1994: 90 (illustration 65)-91, 292 (caption of Table XII), Table XII Fig 1., (5, 6, 9, 265 [as Cantharidae])

Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Middle Miocene (16.0-11.6 Ma). Shanwang Formation.

TYPE LOCALITY. China: Shandong, Shanwang, Linqu County, collection K0.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large mudstone.

COLLECTIONS. Shandong Geological Museum, No. K0349.

COMMENTS AND NOTES. Body length: 11.9 mm x 4.2 mm. Prothorax more large posteriorly, body more short and slender, and last abdominal segment more rounded than the others *Curticantharis*. For this reasons the generic attribution remains doubtful, can be also a species of *Lycocerus*.

†*Themus* sp. (J. Zhang, 1989) [n. comb.]

Curticantharis sp. J. Zhang, 1989: 122, 387, 445 (caption of Table 27), Table 27 Fig. 2., (8 [as Cantharidae])

Zhang et al 1994: 5, 6, 9, 265 [as Cantharidae].

TYPE HORIZON. Middle Miocene (16.0-11.6 Ma). Shanwang Formation.

TYPE LOCALITY. China: Shandong, Shanwang, Linqu County.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large mudstone.

COLLECTIONS. Shandong Provincial Museum (LPM), No. s82753.

COMMENTS AND NOTES. Head and prothorax not preserved. Wing length: 7.8 mm x 2.2 mm. Wing that mostly cover the abdomen.

Genus †*Wongyekokia* Lin, 1997: 186, 187, 188, 189, 194, 199 (caption)

†*Wongyekokia angustris* Lin, 1997

Wongyekokia angustris Lin, 1997: 194 (fig. 6), 199 [caption of the plate 2], pl. 2 fig. 5

Wongyekokia angustis Lee et al., 1997: 17 (incorrect spelling)

Wongyekokia angustis Lin, 1997: 186, 187, 188 (incorrect original spelling. IZN 1999 Art. 19.3)

The Paleobiology Database.

TYPE HORIZON. Paleocene (66-56 Ma). Ping Chau Formation of China.

TYPE LOCALITY. Hong Kong (China), Peng Chau Island (also known as Tung Ping Chau Island), Wong Ye Kok (WYK).

PRESERVATION/LITHOLOGY. Exoskeleton (compression). Lithified, terrestrial siltstone.

COLLECTIONS. Nanjing Institute of Geology and Palaeontology (NIGP), No. 128291 (WYK-4-3).

COMMENTS AND NOTES. Body length: 6.0 mm (preserved only the prothorax, right elytron, and very little part of left elytron), elytron: 4.0 mm x 1.5 mm. For the presence of various striae on elytra, membership in the Cantharidae family is, for me, uncertain. The *angustis* name is earlier to *angustris* but the latter is present in true description (also in the plate and figure), is used in The Paleobiology Database, and particularly there is no evidence to prove it is a *lapsus*.

Tribe Podabrini Gistel, 1856

Genus *Malthacus* Kirby, 1837

***Malthacus deceptus* (W. J. Brown, 1940) [n. comb.]**

Podabrus deceptus W. J. Brown, 1940: 161 (Locus typicus: Churchill, Man., July 13, 1937. Paratypes: June 28 and August 2)

Podabrus deceptus Matthews & Telka, 1997: 921 (Table 1.).

TYPE HORIZON. Early Holocene (9670 ± 130 years old).

TYPE LOCALITY. U.S.A.: northern Alaska, Ikpikpuk River.

PRESERVATION/LITHOLOGY. Fragments or complete body. Ponds, peatland turf.

COLLECTIONS. Geological Survey of Canada, Ottawa.

COMMENTS AND NOTES. Several specimens. *Podabrus deceptus* Brown (1940) was transferred to *Dichelotarsus* by Pelletier & Hébert (2014: 1, 3, 170, 241), but this latter genus *Dichelotarsus* is considered synonymous with *Malthacus* (Takahashi 2007: 241, 243) and therefore the actually correct combination is *Malthacus deceptus* (Brown).

***Malthacus piniphilus* species-group**

Podabrus piniphilus sp. group Matthews & Telka, 1997: 921 (Table 1.)

TYPE HORIZON. Early Holocene (9670 ± 130 years old).

TYPE LOCALITY. U.S.A.: northern Alaska, Ikpikpuk River.

PRESERVATION/LITHOLOGY. Fragments or complete body. Ponds, peatland turf.

COLLECTIONS. Geological Survey of Canada, Ottawa.

COMMENTS AND NOTES. Several specimens.

Genus *Podabrus* Dejean, 1833

***Podabrus* cfr. *alpinus* (Paykull, 1798)**

Podabrus cf. *alpinus* Böcher, 1995: 16, 41-42, 56 (and Table 10.), Table 12. (p. 59), Table 21. [only Cantharidae]

Matthews & Telka 1997: 936 (Table 3.); The Paleobiology Database.

TYPE HORIZON. Pleistocene: Gelasian (about 2 Ma / 1.8-2.2 Ma) - Plio-Pleistocene transition period. Kap København Formation. Late Pliocene (Matthews & Telka 1997: 940 Table 3.).

TYPE LOCALITY. North Greenland (Denmark), eastern Peary Land, Kap København (latitude 82°30'N). Unit B3, locality 77 (Böcher 1995: 9).

PRESERVATION/LITHOLOGY. Insect fragment. Shoreface, poorly lithified siliciclastic sediments.

COLLECTIONS. Zoological Museum, University of Copenhagen (ZMUC).

COMMENTS AND NOTES. One head. The species *Podabrus alpinus* is still alive and lives in a vast territory in Central and Northern Europe from France, Great Britain to Siberia and Mongolia.

†*Podabrus cupesoides* Wickham, 1917

Podabrus cupesoides Wickham, 1917: 463, 467, 472 (explanation of plates), Plate 37 Fig. 4. [under family Lampyridae]

Podabrus cupesoides Wickham, 1913b: 360 (*nomen nudum*) [under family Lampyridae]

Podabrus cupetoides Cossmann, 1918: 13 (incorrect spelling) Cossmann 1918: 13-14; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Podabrus*]; Kirejtshuk & Ponomarenko 2009-2015; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale.

COLLECTIONS. United States National Museum (U.S.N.M.), No. 63446.

COMMENTS AND NOTES. Body length: 8 mm, elytra 5.35 x 2.30 mm. Described from one specimen, but the descriptor associated with this species other specimens, especially those of the Museum of Comparative Zoology and the University of Colorado, that may also be other species. Separated from other Florissant fossil *Podabrus*, for the small size and long antennae reaching nearly to the elytral apex. The assignment to genus *Podabrus* is doubt for Carpenter (1992: 331). The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

†*Podabrus florissantensis* Wickham, 1914

Podabrus florissantensis Wickham, 1914: 444-445, Plate 4. Fig. 2. (and explanation) [under family Lampyridae]

International Catalogue of Scientific Literature 1916b: 147; International Catalogue of Scientific Literature 1919: 108; Meunier 1920a: 156; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Podabrus*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale.

COLLECTIONS. Museum of Comparative Zoology - Harvard University (MCZ), No. 2,495 (No. 8,947 S. H. Scudder coll.).

COMMENTS AND NOTES. Body length: 10.10 mm, elytra 7.35 mm. Described from one specimen. Similar to the fossil *Podabrus fragmentatus* Wickham (see under this species). The assignment to genus *Podabrus* is doubt for Carpenter (1992: 331), and even the descriptor had doubts with the genus *Epicauta* (Coleoptera Meloidae) but distinguishable for the form of the eyes. The picture of the holotype is visible in

Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

†*Podabrus fragmentatus* Wickham, 1914

Podabrus fragmentatus Wickham, 1914: 444, Plate 4. Fig. 1. (and explanation) [under family Lampyridae]

International Catalogue of Scientific Literature 1916b: 147; International Catalogue of Scientific Literature 1919: 108; Meunier 1920a: 156; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Podabrus*]; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale.

COLLECTIONS. Holotype: Museum of Comparative Zoology - Harvard University (MCZ), No. 2,487-2,488 (No. 4,218-4,638 S. H. Scudder coll.). Two other specimens MCZ, No. 2,489-2,490 (No. 69-2,546 S. H. Scudder coll.).

COMMENTS AND NOTES. Body length: 12.75 mm, elytra 6.85 mm. Described from one specimen with counterpart. Similar to the fossil *Podabrus florissantensis* Wickham, but this latter has a larger head, longer elytra and shorter antennal joints (Wickham 1914: 444). The assignment to genus *Podabrus* is doubt for Carpenter (1992: 331). The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

†*Podabrus santaritensis* Cockerell, 1936

Podabrus (?) santaritensis Cockerell, 1936: 2, 5 (and Fig. 5.) Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Paleocene (65.5-56.0 Ma). Maíz Gordo Formation (Sunchal).

TYPE LOCALITY. Argentina: Province of Jujuy, Quebrada "El Griton" (locality oo/cd 27).

PRESERVATION/LITHOLOGY. Elytron (impression). Lacustrine - small mudstone.

COLLECTIONS. American Museum of Natural History, No. 24520, collected by D. C. Harrell.

COMMENTS AND NOTES. One elytron. Length: 4 mm but apex not visible. Very pale brownish, doubtless yellow in life. For the presence of three parallel, widely spaced, strong raised lines and two faint longitudinal rugulose lines between these, is probably a representative of the genus *Podabrus*, but could be also a different but related genus

(Cockerell 1936: 5).

†*Podabrus wheeleri* Wickham, 1909

Podabrus wheeleri Wickham, 1909: 128-129.

Wheeler 1906: figure at the p. 202 (unnamed; "undescribed species of fossil beetle (*Meloid*)"; Wickham 1913a: 284, 293 [under family Lampyridae]; Wickham 1913b: 360 [under family Lampyridae]; Wickham 1914: 444 [under family Lampyridae]; Lucas 1915: 132; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331; White 1995: 37; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant Lake Beds, Station 13.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale.

COLLECTIONS. Holotype in Peabody Museum of Yale University, Cat. No. 7 (Wickham 1909: 129), collection number 165 and received from Prof. Cockerell (Wickham 1909: 128). Two specimens in rather poor condition in United States National Museum (U.S.N.M.), Cat. No. 59653 (Wickham 1913a: 293). Hypotype YPM 7 at Peabody Museum of Natural History Yale University (White 1995: 37 [Wheeler 1906 (p. 202) and Wickham 1909 (p. 128)]). Three specimens according Wickham (1914: 444) preserved at MCZ, No. 2,491, 2,492, 2,493 (No. 5,946, ?, ?, S. H. Scudder coll.) and probably attributable to this species also the specimen No. 2,494 (No. 11,165 S. H. Scudder coll.).

COMMENTS AND NOTES. Body length: 17.25 mm, probably abnormal distended during the fossilization, elytra 11.0, and antenna 7.50 mm. Preserved both the obverse and the reverse. First illustration in Wheeler (1906: fig. at the page 202). The assignment to genus *Podabrus* is doubt for Carpenter (1992: 331). The pictures of the holotype are visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

***Podabrus* spp. Cong, Ashworth, Schwert & Totten, 1996**

Podabrus spp. Cong et al., 1996: 220 (Table 1)

The Paleobiology Database.

TYPE HORIZON. Pleistocene: Illinoian to late Wisconsinan / Tarantian (36.300-43.500 years old). Titusville Till Formation.

TYPE LOCALITY. U.S.A.: Pennsylvania, Titusville Site (41°36'58"N - 79°38'22"W).

PRESERVATION/LITHOLOGY. Pronotum. Mire or swamp, unlithified peat.

COLLECTIONS. Quaternary Entomology Laboratory at North Dakota State University.

COMMENTS AND NOTES. 5 pronotum. One from sample 3 (Zone III: 0-60 cm) and 4 from sample 4 (Zone II: 120-140 cm).

***Podabrus* sp. Matthews & Telka, 1997**

Podabrus sp. Matthews & Telka, 1997: 927 (Table 2.)

TYPE HORIZON. Late Pleistocene: Late Wisconsinan (25460 ± 200 years old).

TYPE LOCALITY. Canada: Yukon, Bell Basin, Rock River exposure.

PRESERVATION/LITHOLOGY. Fragments or complete body. Various sediments.

COLLECTIONS. Uncertain.

COMMENTS AND NOTES. Various specimens. Is also very probable that these specimens are representatives of the genus *Malthacus*.

***Podabrus* sp. Matthews & Telka, 1997**

Podabrus sp. Matthews & Telka, 1997: 936 (Table 3.)

TYPE HORIZON. Pliocene (3-5 Ma).

TYPE LOCALITY. Canada: Ellesmere Island, Strathcona Fiord, Beaver Pond deposit (Beaver peat).

PRESERVATION/LITHOLOGY. Fragments or complete body. Pond.

COLLECTIONS. Uncertain.

COMMENTS AND NOTES. Various specimens.

Subfamily Chauliognathinae LeConte, 1861

Tribe Chauliognathini LeConte, 1861

Genus *Chauliognathus* Hentz, 1830

†*Chauliognathus pristinus* Scudder, 1876

Chauliognathus pristinus Scudder, 1876: 81

Chauliognathus Handlirsch, 1920-1921: 232 (incorrect spelling)

Goss 1878: 63 (reprint 1878: 341); Scudder 1882: 292; Scudder 1885: 796 (Fig. 1032.) [under family Lampyridae]; Scudder 1886: 76 [under family Lampyridae]; Scudder 1887: 796 (Fig. 1050.) [under family Lampyridae]; Scudder 1891: 489 [under family Lampyridae]; Scudder 1900: 101-102, Plate XI Fig. 3. [under family Lampyridae]; Handlirsch 1906-1908: 739; Grabau & Shimer 1910: 447, 708 (Appendix B.) [as fire-fly]; Wickham 1913a: 284, 292-293 [under family Lampyridae]; Lucas 1915: 134 [under family Lampyridae]; Wickham 1920: 353; Handlirsch 1920-1921: 232; Miskimen 1961: 149-152; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331; Douglas & Stockey 1996: 1154; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant, Castello's ranch.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale.

COLLECTIONS. United States National Museum (U.S.N.M.), Cat. No. 59652, for the specimen in Lacoe collection (Wickham 1913a: 293).

COMMENTS AND NOTES. Redescribed by Scudder (1900: 101-102). Body length: 13.0 mm (elytron 6.0 mm), antennae 6.0 mm. Specimen in Lacoe collection: body length 10.60 mm and elytra 6.0 mm (Wickham 1913a: 293). The assignment to genus *Chauliognathus* is doubt for Carpenter (1992: 331), and would have confirmed by the fact that in the original description, Scudder (1876: 81) and the same author (Scudder 1900: 101) says apparently having 12 antennomeres. Drawning in Scudder (1885: fig. 1032; 1887: fig. 1050; 1900: plate XI, fig. 3 [drawning by Paul Roetter]), T. L. Mead *legit*). The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

Tribe Ichthyurini Champion, 1915

Genus *Trypherus* LeConte, 1851

†*Trypherus aboriginalis* Wickham, 1913

Trypherus aboriginalis Wickham, 1913b: 360, 362-363, 366 (explanation of plates), Plate XXXVIII. Fig. 3. (but indicated as Plate I. Fig. 3. in the text and explanation) [under family Lampyridae]

Wickham 1914: 446 [under family Lampyridae]; Lucas 1915: 132; International Catalogue of Scientific Literature 1916a: 95; Meunier 1920b: 159; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Trypherus*]; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database [under family Lampyridae]; The Paleobiology Database.

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale, volcaniclastic.

COLLECTIONS. Holotype: Museum of Princeton University, No. 6527. And two specimens, one with counterpart, preserved at MCZ, No. 2,503-2,505 (No. 8,586, 8,499, 8,651 S. H. Scudder coll.).

COMMENTS AND NOTES. Body length: 8.75 mm. Similar to the recent *Trypherus latipennis* (Germar). The assignment to genus *Trypherus* is doubt for Carpenter (1992: 331). The picture of the holotype is visible in Florissant Fossil Beds National Monument website at <http://planning.nps.gov/flfo/photos>.

Subfamily Dysmorphocerinae Brancucci, 1980

Genus †*Cacomorphocerus* Schaufuss, 1892: 57-58

= *Hoffeinsensia* Kuška & Kania, 2010: 50 [synonymized by Kazantsev, 2013: 289, 290]

Hoffeinsensis Pielińska, 2006: 9, english part: 34 (incorrect spelling)

†*Cacomorphocerus cerambyx* Schaufuss, 1892

Cacomorphocerus cerambyx Schaufuss, 1892: 58

Carus (ed.) 1892: 84; Helm 1897: 88-89; Schröder 1900: 355; Handlirsch 1906-1908: 740; Handlirsch 1920-1921: 232; Korschefsky 1939: 11-12, Tafel 1 (figs. 2abc); Weidner 1952: 69 [only Cantharidae], 71 [*Cacomorphocerus* ähnlich *cerambyx*]; Larsson 1978: 140; Spahr 1981b: 15; Winkler 1987: 58; Carpenter 1992: 306; Poinar 1992: 137; Kuška 1994: 6; Douglas & Stockey 1996: 1154; Lin 1997: 189; Hoffeins 2008: 39, 40; Alekseev 2013: 8; Kazantsev 2013: 283, 289, 290; Kazantsev & Perkovsky 2014: 116; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Westpreußischen Provinzial-Museum Danzig (= Museum für Naturkunde und Vorgeschichte, WPM), Coll. Helm, No. 67.

COMMENTS AND NOTES. Sex undefined. Body length: 6 mm x 1,75-2 mm. Genus similar to the extant *Dysmorphocerus* Solier in Gay, characterized to antennae 12-segmented with articles 3-9 dilated and saucer-shaped. Drawnings of type not printed by Schaufuss (1892) can be found in Korschefsky (1939: table 1. Drawnings of Mühling), table also then taken up by Hoffeins (2008: 40). Korschefsky (1939: 11) cited, and the data was taken by Weidner (1952: 71), another specimen related to *Cacomorphocerus cerambyx* in Scheele (Berlin/Lichterfelde) collection at University of Hamburg from Eocene of Russian: Kaliningrad region (The Paleobiology Database) but with 16 (or 15, because the text is not clear) antennomeres. And for these reasons I believe that you can treat of a genus (surely a new species) not described yet. Moreover for Winkler (1987: 58) the genus *Cacomorphocerus* is not a true Cantharid, but probably has affinities with Phengodidae or Teleugesidae.

†*Cacomorphocerus jantaricus* (Kuška & Kania,

2010)

Hoffeinsensia jantarica Kuška & Kania, 2010: 50-52 (figs. 1-2 p. 51 and figs. 3-6 p. 52)
Pawlowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 289-290; Kazantsev & Perkovsky 2014: 116; Kania 2015: 12; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber (East Baltic Amber).

COLLECTIONS. Deutsches Entomologisches Institut (DEI) in Germany, No. 1221-2 and 1226-1 (ex coll. C. and H. W. Hoffeins, Hamburg). Three specimens not in type series in private collection of Jacek Serafin (Piaseczno- Poland).

COMMENTS AND NOTES. 4 males, and 1 probably it is also a male. Pronotum subrectangular brown with two lateral protuberances in the basal half, and elytra almost black. Body length: holotype 6.5 mm, others: 7.0-9.0 mm. Various syninclusions. Transferred to *Cacomorphocerus* by Kazantsev (2013: 289-290).

†*Cacomorphocerus* sp. Kazantsev & Perkovsky, 2014

Cacomorphocerus sp. Kazantsev & Perkovsky 2014: 116

TYPE HORIZON. Upper Eocene: Priabonian (38.0/37.2-33.9 Ma) or Lower Oligocene.

TYPE LOCALITY. Ukraine: Klesov, Pugach quarry.

PRESERVATION/LITHOLOGY. Inclusion in amber.

Rovno Amber.

COLLECTIONS. Schmalhausen Institute of Zoology of Kiev, No. K-541.

COMMENTS AND NOTES. One male. Syninclusions: One Diptera (Chironomidae).

Subfamily Malthininae Kiesenwetter, 1852

Tribe Malchinini Brancucci, 1980

Genus *Macrocerus* Motschulsky, 1845

†*Macrocerus sucinopenninus* (Kuška & Kania, 2010)

Malthodes sucinopenninus Kuška & Kania, 2010: 54-55 (figs. 14-16)

Pawlowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-

33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Deutsches Entomologisches Institut (DEI) in Germany, No. 1221 -1 (ex coll. C. and H. W. Hoffeins, Hamburg).

COMMENTS AND NOTES. Male, brownish-black. Body length: 2.4 mm. Similar to the extant *Malthodes penninus* Baudi di Selve. Syninclusions: *exuvium* of Orthoptera and two tufts of oak hairs. Transferred to *Macrocerus* by Kazantsev (2013: 290).

†*Macrocerus* sp. (Klebs, 1910)

Malchinus Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Spahr 1981b: 15; Hieke & Pietrzeniuk 1984: 303, 321; Poinar & Fanti 2016: 1.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection, collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Three specimens.

†*Macrocerus* sp. (Bachofen-Echt, 1949)

Malchinus Bachofen-Echt, 1949: 109

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Macrocerus* sp. (Kubisz, 2000)

Malchinus Kubisz, 2000: 226

Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland ?: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of Natural History of the Institute of Systematics and Evolution of Animals (ISEA) in Kraków.

COMMENTS AND NOTES. One specimen.

Tribe Malthinini Kiesenwetter, 1852

Genus *Malthinus* Latreille, 1806

Subgenus (*Malthinus*) Latreille, 1806

†*Malthinus (Malthinus) danieli* Kuška & Kania, 2010

Malthinus danieli Kuška & Kania, 2010: 53-54 (figs. 10-13); Pawłowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Deutsches Entomologisches Institut (DEI) in Germany, No. 161-1 (ex coll. C. and H. W. Hoffeins, Hamburg).

COMMENTS AND NOTES. Male, greyish-brown with pronotum yellow with black spot on posterior half, tibiae yellow-brown and femora dark. Body length: 3.2 mm, antennae 3.5 mm long. Similar to the extant *Malthinus seriepunctatus* Kiesenwetter.

†*Malthinus* sp. Hope, 1836

Malthinus Hope, 1836: 143 (Berendt authority).

Berendt 1845: 56; Pictet 1854: 334; Giebel 1856a: 100-101 [under family Lampyridae], 400 (in the "Tabellarische"); Giebel, 1856b: 181; Oustalet 1874: 22 [citation of Pictet]; Scudder 1891: 550 [under family Lampyridae]; Handlirsch 1906-1908: 740 [as *Malthinus* - Berendt]; Larsson 1978: 139; Spahr 1981b: 4, 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber. (Prussian Amber).

COLLECTIONS. Berlin Museum (coll. Berendt).

COMMENTS AND NOTES. One specimen.

†*Malthinus* sp. Giebel, 1852

Malthinus Giebel, 1852: 655

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. One specimen.

†*Malthinus* sp. Motschulsky, 1857

Malthinus Motschulsky, 1857: 28

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Coll. Menge.

COMMENTS AND NOTES. One or more specimens.

†*Malthinus* sp. Scudder, 1885

Malthinus Scudder, 1885: 796 [under family Lampyridae]

Scudder 1886: 76 [under family Lampyridae]; Scudder 1887: 796 [under family Lampyridae]; Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. One specimen, and probably the same as some previous author (probably Berendt, 1845).

†*Malthinus* sp. Helm, 1896

Malthinus Helm, 1896: 228

Handlirsch 1906-1908: 741; Handlirsch 1920-1921: 232; Larsson 1978: 139; Spahr 1981b: 15; Kazantsev 2013: 283.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Malthinus* sp. Klebs, 1910

Malthinus Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 15, 16; Hieke & Pietreniuk 1984: 303, 321; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection, collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Four specimens, and one near to *Malthinus*.

†*Malthinus* sp. Bachofen-Echt, 1949

Malthinus Bachofen-Echt, 1949: 109

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Malthinus* sp. Hieke & Pietrzeniuk, 1984

Malthinus Hieke & Pietrzeniuk, 1984: 303

Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museums für Naturkunde, Berlin.

COMMENTS AND NOTES. Hieke & Pietrzeniuk (1984: 303, 321) quote the genera: *Cantharis*, *Rhagonycha*, *Malthinus* and *Malthodes*, studying about 110 specimens of various Museums and collections: Berlin: Künow collection - 25 specimens, Berendt collection - 10 specimens, Kühl collection - 7 specimens, Simon collection - 3 specimens, Isenheim collection - 3 specimens, "Bitterfelder Bernstein" - 1 specimen, Mineralogischen Museum - 1 specimen; Klebs collection - 47 specimens; Copenhagen collection - ? few.

Tribe Malthodini Böving & Craighead, 1931

Genus *Archaeomalthodes* Hsiao, Ślipiński & Pang, 2016: 121

†*Archaeomalthodes rosetta* Hsiao, Ślipiński & Pang, 2016

Hsiao et al., 2016: 121-122 (figs. 2-4), 123

Fanti & Ellenberger 2016: 166, 169; The Paleobiology Database.

TYPE HORIZON. Early Cretaceous: Upper Albian (97-110 Ma).

TYPE LOCALITY. Myanmar: Kachin State, Hukawng Valley, approximately 100 km southwest of the Village of Tanai.

PRESERVATION/LITHOLOGY. Inclusion in amber. Burmese Amber.

COLLECTIONS. Key Laboratory of Insect Evolution & Environmental Changes, Capital Normal University, Beijing (China), CNU-COL-MA20160200.

COMMENTS AND NOTES. Male. Brown, with black eyes, and covered with fine yellowish pubescence. Body length: 2.50 mm (from the anterior margin of the clypeus to the apices of elytra). Elytra somewhat abbreviated with caudal abdominal segments exposed (Hsiao et al. 2016: 121).

Genus *Malthodes* Kiesenwetter, 1852

Subgenus (*Malthodes*) Kiesenwetter, 1852

†*Malthodes* (*Malthodes*) *ceranowiczae* Kuśka & Kupryjanowicz, 2005

Malthodes ceranowiczae Kuśka & Kupryjanowicz, 2005: 310-311 (figs. 1-4), 314 (figs. 9-10)

Malthodes ceranoviczae Kazantsev, 2013: 290 (incorrect spelling)

Malthodes ceranoviczae Kazantsev & Perkovsky, 2014: 114 (incorrect spelling)

Kuśka & Kania 2010: 55; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113, 114; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of the Earth PAS in Warsaw (Museum Ziemi collection), No. 24198.

COMMENTS AND NOTES. Two males in one piece of amber. Brown with head and pronotum slightly darker (brown-black). Body length: 2.7 mm, elytra 1.28 mm, width of elytra 0.48 mm, width of pronotum at the base 0.4 mm, antennae 2.0 mm. Related to the extant species group *M. maurus* and particularly to *M. caudatus* Weise. Similar specimen of *M. ceranowiczae* is present in private collection to Andrzej Górski at Bielsko-Biala (Kuśka & Kupryjanowicz 2005: 311).

†*Malthodes* (*Malthodes*) *kotejai* Kuśka & Kupryjanowicz, 2005

FOSSILS & MINERALS

Malthodes kotejai Kuška & Kupryjanowicz, 2005: 311-312 (figs. 5-7), 315 (fig. 11)

Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113, 114; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of the Earth PAS in Warsaw (Museum Ziemi collection), No. 24199. (coll. J. Kupryjanowicz).

COMMENTS AND NOTES. Male, brown with pronotum almost yellow and head dark brown. Body length: 3.4 mm, elytra 1.37 mm, width of elytra 0.65 mm, antennae 3.25 mm. Terminalia very simple.

†*Malthodes (Malthodes) obtusus* Förster, 1891

Malthodes obtusus Förster, 1891: 373-374, Tafel XI: Fig. 18

Scudder 1891: 550 [only as *Malthodes* Foerster in litt. (under family Lampyridae)]; Handlirsch 1906-1908: 741; Handlirsch 1920-1921: 232; Théobald 1937: Tableau X, 273; Carpenter 1992: 331; Kazantsev 2010: 105; Kazantsev & Perkovsky 2014: 113, 115; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Early/Lower Oligocene (33.9-28.4 Ma) sometimes considered Eocene: Priabonian.

TYPE LOCALITY. France: Alsace, Brunstatt (horizon d3).

PRESERVATION/LITHOLOGY. Mold impression/compression (adpression). Lacustrine lithified marl.

COLLECTIONS. Not indicated, and probably lost or destroyed during World War II.

COMMENTS AND NOTES. Sex undefined. Light brown. Body length: around 3.6 mm (because elytra 2.1 mm, head 0.6 mm, and prothorax 0.6/0.9 mm). Terminalia not visible. The assignment to genus *Malthodes* is doubt for Carpenter (1992: 331).

†*Malthodes (Malthodes) perkovskyi* Kazantsev, 2010

Malthodes perkovskyi Kazantsev, 2010: 105-107 (figs. 1-3 pp. 106-107)

Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113, 114, 115; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Upper Eocene: Priabonian (38.0/37.2-33.9 Ma) or Lower Oligocene.

TYPE LOCALITY. Ukraine: Klesov.

PRESERVATION/LITHOLOGY. Inclusion in amber. Rovno Amber.

COLLECTIONS. Schmalhausen Institute of Zoology

of Kiev, No. K-4579.

COMMENTS AND NOTES. Male, dark brown to black, pronotum lighter with black W-sign, elytra with apically yellow spots. Body length: 3.4 mm, width 0.6 mm. Related to the extant Japanese and Southern Kurils: *Malthodes kurosawai* Wittmer and *Malthodes kerzhneri* Wittmer.

†*Malthodes (Malthodes) rovnoensis* Kazantsev & Perkovsky, 2014

Malthodes rovnoensis Kazantsev & Perkovsky, 2014: 113-114 (figs. 1-2), figs. 3-6 (p. 115)

Perkovsky 2015: 410; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; The Paleobiology Database.

TYPE HORIZON. Upper Eocene: Priabonian (38.0/37.2-33.9 Ma) or Lower Oligocene.

TYPE LOCALITY. Ukraine: Klesov, Pugach quarry.

PRESERVATION/LITHOLOGY. Inclusion in amber. Rovno Amber.

COLLECTIONS. Schmalhausen Institute of Zoology of Kiev, No. K-25818.

COMMENTS AND NOTES. Male, brownish with head black. Body length: ca. 2.5 mm and width ca. 0.6 mm. Similar to the extant *M. minimus* Linnaeus, and fossil *M. perkovskyi* Kazantsev.

†*Malthodes (Malthodes) serafini* Kuška & Kupryjanowicz, 2005

Malthodes serafini Kuška & Kupryjanowicz, 2005: 312-313 (fig. 8), 315 (fig. 12)

Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113, 115; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Natural History Museum, ISEA of Krakow, No. MP/1/1513/188/01.

COMMENTS AND NOTES. Male, black. Body length: 1.9 mm, elytra 1.25 and width 0.56 mm, width of antennae 2.0 mm. Similar to the extant *M. hexacanthus* Kiesenwetter. Syninclusions: 4 specimens of Chironomidae and 1 of Acarina.

†*Malthodes (Malthodes) sucini* Kuška & Kania, 2010

Malthodes sucini Kuška & Kania, 2010: 55-56 (figs. 17-18) Pawłowski & Mazur 2012: 44; Alekseev 2013: 8; Kazantsev 2013: 283, 290; Kazantsev & Perkovsky 2014: 113, 115; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; Mitchell 2013 - EDNA

The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Deutsches Entomologisches Institut (DEI) in Germany, No. 161-2 (ex coll. C. and H. W. Hoffeins, Hamburg).

COMMENTS AND NOTES. Male. Body length: 2.8 mm, elytra 1.4 mm long. Similar to the fossil *Malthodes ceranowiczae* Kuška & Kupryjanowicz that differs for the last sternite shape.

†*Malthodes* sp. Klebs, 1910

Malthodes Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 16; Hieke & Pietrzeniuk 1984: 303, 321; Kazantsev 2010: 105; Kazantsev 2013: 283; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection, collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Five specimens.

†*Malthodes* sp. Bachofen-Echt, 1949

Malthodes (sic!) Bachofen-Echt, 1949: 109

Spahr 1981b: 16.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

†*Malthodes* sp. Hieke & Pietrzeniuk, 1984

Malthodes Hieke & Pietrzeniuk, 1984: 303

Kazantsev 2010: 105; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museums für Naturkunde, Berlin.

COMMENTS AND NOTES. Hieke & Pietrzeniuk (1984: 303, 321) quote the genera: *Cantharis*, *Rhagonycha*, *Malthinus* and *Malthodes*, studying about 110 specimens of various Museums and collections: Berlin: Künnow collection - 25 specimens, Berendt collection - 10 specimens, Kühl collection - 7 specimens, Simon collection - 3 specimens, Isenheim collection - 3 specimens, "Bitterfelder Bernstein" - 1 specimen, Mineralogischen Museum - 1 specimen; Klebs collection - 47 specimens; Copenhagen collection - ? few.

†*Malthodes* sp. Heynderycx, 2004

Malthodes sp. Heynderycx, 2004: 3

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. ?

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Thadeus Giecewicza collection (Warsaw - Poland).

COMMENTS AND NOTES. One specimen.

†*Malthodes* sp. Penney & Jepson, 2014

Malthodes sp. Penney & Jepson, 2014: Fig. 215 (p. 173)

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Not indicated

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. RCDP: Research collection David Penney, UK.

COMMENTS AND NOTES. One specimen with wing extended.

Tribe †*Mimoplatycini* Kazantsev, 2013: 288,

290

Genus †*Mimoplatycis* Kazantsev, 2013: 288,

290

†*Mimoplatycis notha* Kazantsev, 2013

Mimoplatycis notha Kazantsev, 2013: 288 (figs. 10-11)-289 (figs. 12-14), 290

Kazantsev & Perkovsky 2014: 116 (figs. 7-9); Hsiao et al. 2016: 119, 120, 122; Poinar & Fanti 2016: 1, 2; The Paleobiology Database.

FOSSILS & MINERALS

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma). Rovno amber: Upper Eocene: Priabonian (38.0/37.2-33.9 Ma) or Lower Oligocene.

TYPE LOCALITY. Denmark: Baltic Sea coast. Paratype of Russian Federation: Kaliningrad region. Known also for Ukraine: Klesov, Pugach quarry (Kazantsev & Perkovsky 2014: 113, 116).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber - Scandinavian Amber (Paratype: East Baltic Amber). Known also for Rovno Amber.

COLLECTIONS. Zoological Museum of Copenhagen University, No. 1-12/1966. Paratype in private collection of V. I. Alekseev (Kaliningrad - Russia), No. AWI-038. Specimens from Rovno amber (2 males) collected at the Schmalhausen Institute of Zoology of Kiev with numbers: K-6089 and UA-2286.

COMMENTS AND NOTES. Two males. Holotype: dark brown with pronotal sides and elytra yellow-brown, and only apex of elytra darkened. Paratype: more darkened, with elytra dark-brown. Body length: 2.9-3.6 mm, width 0.7-0.9 mm. Characterized for six male ventrites and carinate pronotum. Other two males in Rovno amber.

Subfamily Silinae Mulsant, 1862

Tribe Silini Mulsant, 1862

Genus *Autosilis* Kazantsev, 2011

cfr. *Autosilis nitidula* (Fabricius, 1792)
Spilis (sic!) *spinicolis* (sic!) Serres, 1843: 35
Pictet 1854: 335 [as *Silis spinicollis*].

TYPE HORIZON. Tertiary strata (Eocene). (Eocene-Oligocene?).

TYPE LOCALITY. France: Aix-en-Provence.

PRESERVATION/LITHOLOGY. Argillaceous limestone and marl.

COLLECTIONS. ?

COMMENTS AND NOTES. Probably one fragment. Specimen of this species or one very similar.

Genus †*Curche* Alekseev & Kazantsev, 2014: 167-168

†*Curche pauli* Alekseev & Kazantsev, 2014

Curche pauli Alekseev & Kazantsev, 2014: 167-170 (figs. 1-5 p. 169)

Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4

Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Kaliningrad region, Baltic Sea coast, Yantarny settlement (formerly Palmnicken).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Paleontological Institute, Russian Academy of Science (Moscow), No. AWI-100, ex coll. Alekseev (Kaliningrad - Russia).

COMMENTS AND NOTES. Male, black with pronotum, legs and head appendages reddish-brown (in lifetime probably red or orange). Body length: 7.2 mm, width 2.5 mm. Syninclusions: a spider 3.7 mm in length.

Genus †*Electrosilis* Kazantsev, 2013: 285, 290

†*Electrosilis minuta* Kazantsev, 2013

Electrosilis minuta Kazantsev, 2013: 285-286 (figs. 4-6), 290 Alekseev & Kazantsev 2014: 167, 168; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; The Paleobiology Database.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Russian Federation: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Senckenberg Deutsches Entomologisches Institut (Müncheberg - Germany), No. 1441-2.

COMMENTS AND NOTES. Male, light brown. Body length: 2.1 mm, width 0.7 mm. The genus *Electrosilis* is similar to *Silis* Charpentier, and distinguishable by the narrowly explanate pronotal sides.

Genus *Polemius* LeConte, 1851

†*Polemius crassicornis* Wickham, 1914

Polemius crassicornis Wickham, 1914: 445-446, Plate 4. Fig. 4. (and explanation) [under family Lampyridae]

International Catalogue of Scientific Literature 1916b: 147; International Catalogue of Scientific Literature 1919: 108; Meunier 1920a: 156; Wickham 1920: 353; Handlirsch 1920-1921: 232; Wilson 1978: 21 (Table 1.); Carpenter 1992: 331 [as *Polemius*]; Kirejtshuk & Ponomarenko 2009-2015 [as ?*Polemius* ???????]; Mitchell 2013 - EDNA The Fossil Insect Database [under family Lampyridae]; The Paleobiology Database [under family Lampyridae].

TYPE HORIZON. Eocene: Chadronian (37.2-33.9 Ma), Florissant Formation.

TYPE LOCALITY. U.S.A.: Colorado, Teller County, Florissant.

PRESERVATION/LITHOLOGY. Compression/

adpression. Lacustrine - large shale, diatomaceous and tuffaceous shale, volcaniclastic.

COLLECTIONS. Museum of Comparative Zoology - Harvard University (MCZ), No. 2,502 (No. 930 S. H. Scudder coll.).

COMMENTS AND NOTES. Head nearly concealed. Body length: 8.40 mm. Described from one specimen. The assignment to genus *Polemius* is doubt for Carpenter (1992: 331).

Genus *Silis* Charpentier, 1825

†*Silis chiapasensis* Wittmer, 1963

Silis chiapasensis Wittmer, 1963: 53 (fig. 136), Plate 3: lower right (p. 59)

Hurd et al. 1962: 110 [only as Cantharidae, but probably this species]; Spahr 1981b: 15 [Hurd], 16; Perrilliat 1989: 394 (Figura 153. f), 396; Poinar 1992: 137 (Figure 76); Kuška 1994: 6; Douglas & Stockey 1996: 1154; Engel 2004: 184; Cifuentes-Ruiz et al. 2007: 678; Solórzano Kraemer 2007: 120; Solórzano Kraemer 2010: 56; Kazantsev 2013: 283, 290; Alekseev & Kazantsev 2014: 167; Hsiao et al. 2016: 120; Poinar & Fanti 2016: 1; [authorless]: 39; Kirejtshuk & Ponomarenko 2009-2015; Mitchell 2013 - EDNA The Fossil Insect Database; The Paleobiology Database.

TYPE HORIZON. Early/Lower Miocene (23.0-16.0 Ma) or Middle Miocene (Solórzano Kraemer 2007) sometimes considered also Late Oligocene.

TYPE LOCALITY. Mexico, Chiapas, Simojovel de Allende area (locality B-1402).

PRESERVATION/LITHOLOGY. Inclusion in amber. Mexican (Chiapas) Amber.

COLLECTIONS. University of California Museum of Paleontology (UCMP), No. 12615.

COMMENTS AND NOTES. Male, yellow-brown, elytra dark. Body length: 3.5 mm to apex of elytra but 4.5 mm with protruding abdominal segments included. Pronotum with deeply incision in front of the anterior angles and this with pilose process. Photo of the Holotype is present also in Perrilliat (1989: 394 fig. 153f) and Poinar (1992: 137).

†*Silis* sp. Klebs, 1910

Silis Klebs, 1910: 237

Handlirsch 1920-1921: 232; Crowson et al. 1967: 527; Crowson 1972: 63-64 [only as Cantharidae]; Larsson 1978: 140; Spahr 1981b: 16; Hieke & Pietrzeniuk 1984: 303, 321; Kazantsev 2013: 283; Alekseev & Kazantsev 2014: 167; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably, Eastern Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. The vast Klebs's amber collection,

collected in Königsberg in Pr. Museum, has been largely destroyed or stolen during the World War II, and now preserved in several collections, especially in Göttingen University. Hieke & Pietrzeniuk (1984: 300, 303, 321) have seen 47 samples of Cantharidae stored at Halle/Saale and London.

COMMENTS AND NOTES. Two specimens.

†*Silis* sp. Bachofen-Echt, 1949

Silis Bachofen-Echt, 1949: 109

Spahr 1981b: 16.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

Tribe *Tytthonyxini* Arnett, 1962

Genus *Tytthonyx* LeConte, 1851

Subgenus (*Tytthonyx*) LeConte, 1851

†*Tytthonyx* (*Tytthonyx*) *geiseri* Poinar & Fanti, 2016

Tytthonyx geiseri Poinar & Fanti, 2016: 5 (figs. 7-8), 6
Hsiao et al. 2016: 120; The Paleobiology Database.

TYPE HORIZON. Uncertain. Upper Eocene to Lower Miocene (45-30 Ma to 20-15 Ma).

TYPE LOCALITY. Dominican Republic: Cordillera Septentrional, mine between Puerto Plata and Santiago.

PRESERVATION/LITHOLOGY. Inclusion in amber. Dominican Amber.

COLLECTIONS. George Poinar amber collection at Oregon State University, Corvallis (USA), No. C-7-70.

COMMENTS AND NOTES. Male. Body length: 2.4 mm but with terminal abdominal segments folded up (estimated body length: 2.8-3.2 mm). First fossil *Tytthonyx* described and characterized by the small size. Syninclusions: One Diptera.

Specimen undescribeds (family and subfamilies level)

Cantharidae Bachofen-Echt, 1949

Cantharidae Bachofen-Echt, 1949: Abb. 85. (p. 100), p. 102, 108 [also as *Cantharinae*]-109

Crowson 1972: 63-64; Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4

Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber.

Baltic Amber.

COMMENTS AND NOTES. One specimen.

Cantharidae Larsson, 1965

Cantharidae Larsson, 1965: 141

Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Zoological Museum, Copenhagen.

COMMENTS AND NOTES. Two specimens.

Cantharidae Gersdorf, 1971

indeterminiert, vielleicht *Cantharidae* Gersdorf, 1971: Tafel 62: Fig. 4. (pp. 668-669)

The Paleobiology Database.

TYPE HORIZON. Pliocene: Piacenzian (3.6-2.6 Ma).

TYPE LOCALITY. Germany, Niedersachsen, 3411 Willershausen clay pit (Kreis Osterode am Harz).

PRESERVATION/LITHOLOGY. Mold impression/adpression. Pond/little lake, lithified marl and sediments.

COLLECTIONS. GZG collection?. Probably in local Museum.

COMMENTS AND NOTES. Body length: 8 mm. Fairly well preserved. Picture of Riek of Hannover (Gersdorf 1971: T. 62 at pag. 668).

Cantharidae Jell & Duncan, 1986

Cantharid indet. Jell & Duncan, 1986: 114 (Table 1.), 156-157, Fig. 39. A (p. 158), Fig. 40. A (p. 159)

Jell 2004: 72 (and figure and drawing); Nicholson 2012: 271; The Paleobiology Database.

TYPE HORIZON. Lower Cretaceous: Late/Upper Aptian (122.5-112.0 Ma). Strzelecki Group.

TYPE LOCALITY. Australia: Victoria, South Gippsland, Koonwarra Fossil Bed (Site NMVPL425).

PRESERVATION/LITHOLOGY. Adpression. Pond, lithified claystone and lithified siltstone.

COLLECTIONS. Palaeontological Collection, Museum of Victoria, Melbourne (NMVP), No. NMVP103331.

COMMENTS AND NOTES. One adult specimen. Body length: 7 mm. Head narrow with clypeal region produced forward, pronotum much wider than head with lateral margin concave anteriorly, elytra soft not covering apical segments of abdomen, femur

expanded and long thin tibia. May be compared with the primitive living *Telephorus* = *Cantharis* and *Heteromastix* (Jell & Duncan 1986: 157). Refigured and re-drawning in Jell (2004: 72).

Cantharidae Henwood, 1992

Cantharidae Henwood, 1992: 904-908, Plate 1: figs. 1-2, 4-8 (p. 907)

TYPE HORIZON. Uncertain. Upper Eocene to Lower Miocene (45-30 Ma to 20-15 Ma).

TYPE LOCALITY. Dominican Republic, probably Palo Alto in the Cordillera Septentrional.

PRESERVATION/LITHOLOGY. Inclusion in amber. Dominican Amber.

COLLECTIONS. Deposited at Sedgwick Museum, Cambridge, No. SM X.23256 and SM X.23257.

COMMENTS AND NOTES. Body length: about 5 mm. Specimen serially sectioned for the soft-part vision.

Cantharidae? Adamonis, 1994

Cantharidae? Adamonis, 1994: 48

Petrilevicius 1999: 96.

TYPE HORIZON. Eocene: Late Paleocene - Middle Eocene, sometimes considered Oligocene. Formación Ventana.

TYPE LOCALITY. Argentina: Neuquén province (border with Rio Negro province), Confluencia Department, confluence of the River Limay with River Traful.

Cantharidae Douglas & Stockey, 1996

Cantharidae Douglas & Stockey, 1996: 1142, Fig. 7. (p. 1144), 1145 (caption of Fig. 7.), 1153, 1154, 1155 (as Cantheridae sic!), 1156

TYPE HORIZON. Middle Eocene (48-50 Ma).

TYPE LOCALITY. Canada: British Columbia, 8 Km north of Princeton, One Mile Creek.

PRESERVATION/LITHOLOGY. Compression/impression. Freshwater lake, silt, clays and finely graded volcanics.

COLLECTIONS. Paleontology collections of the University of Alberta, Edmonton (UAPAL), No. 4513.

COMMENTS AND NOTES. Well preserved but no totally the head and appendages. Body length: 11.7 mm, width 3.5 mm, antenna filiform 3.3 mm (not complete). Femora light brown and oval, tibiae and tarsi slender. Elytra have squared posterior margins, parallel-sides and is melanic. Pronotum light-brown, square, narrower than the elytra. Head broad and prominent, slightly narrower than the pronotum with probably forward-protruding clypeus (Douglas & Stockey 1996: 1142).

Cantharidae Elias, Short & Waythomas, 1996
Cantharidae Genus et sp. indet. Elias, Short & Waythomas, 1996: 298

TYPE HORIZON. Pleistocene/Holocene: mid-Wisconsin - Boutellier interval (60.000-40.000 years old) with other sediments of 8.075-6.540 years old or more recent. Probably this specimen is more recent (Holocene).

TYPE LOCALITY. U.S.A.: Alaska, Denali National Park and Preserve, Foraker River, Foraker Slump (Sample D).

PRESERVATION/LITHOLOGY. Silt - organic silt and peat.

COMMENTS AND NOTES. One specimen.

Thanks to the courteous information of Vinicius S. Ferreira two photos of two different specimens of Cantharidae from Dominican amber are also present in: Wu R.J.C., 1997. Secrets of a Lost World: Dominican Amber and its Inclusions. Privately published, Santo Domingo, Dominican Republic.

Cantharidae Weitschat & Wichard, 1998
Cantharidae Weitschat & Wichard, 1998: 162, 164 (caption of the Tafel 62), 165 (Tafel 62 fig. f), 166
 Weitschat & Wichard 2002: 160, 162 (caption of the Plate 62), 163 (Plate 62 fig. f), 164; Klausnitzer 2003: 54; Binder 2008: 47.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. One specimen.

Cantharidae Janzen, 2002
Cantharidae Janzen, 2002: 111 (Fig. 227), 56
 Klausnitzer 2003: 55.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Coll. Glink.

COMMENTS AND NOTES. Body length: 3.5 mm.

Cantharidae Wichard & Weitschat, 2004
Cantharidae Wichard & Weitschat, 2004: 79 [figure and drawing]
 Arillo 2007: 160; Weitschat 2009: Fig. 21 (p. 248); Wichard & Greven 2009: Abb. 5ab (p. 286).

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. Cantharidae in mating.

Cantharidae Poinar, Marshall & Buckley, 2007 see under *Ornatomalthinus elvira*

Cantharidae Poinar et al., 2007: 1663-1668
 Binder 2008: 47 (Abb. 1); Boucot & Poinar 2010: 196 (Figure 269); Penney & Jepson 2014: 170; Hsiao et al. 2016: 122; Poinar & Fanti 2016: 1.

Cantharidae Wilson, 2008-2009

Cantharidae Wilson, 2008-2009: 29 (Appendix 2), 44 (Appendix 5) [as Cantharidae? Archibald (personal communication, June 2007)]

Bruce Archibald et al. 2010: Online Supplementary Material p. 15 (Online Table 4: Specimen / species compositions of samples) [as cf. Cantharidae].

TYPE HORIZON. Early Eocene (52.90 ± 0.83 Ma). Kamloops Group.

TYPE LOCALITY. Canada: British Columbia, 15 Km east of Cache Creak, McAbee Fossil Beds.

COLLECTIONS. Unknown. Number 2400 (Bruce Archibald et al. 2010: Online Suppl. Mat. p. 15).

PRESERVATION/LITHOLOGY. Fragment?. Lacustrine shale.

COMMENTS AND NOTES. Probably only one specimen.

Cantharidae Wichard, 2009

Cantharidae Wichard, 2009: Abb. 2a-c (p. 260), 261

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. Specimen with syninclusions: Diptera Psychodidae (*Trichomyia*) and Nevorthidae (*Rophalis relicta*).

Cantharidae Wichard & Greven, 2009

Cantharidae Wichard & Greven, 2009: Abb. 5abcd [figures 5ab are the same of Wichard & Weitschat 2004: 79], 286

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. Three specimens (three species) with one with two exemplars in mating (the same of Wichard & Weitschat 2004: 79).

Cantharidae Wedmann, Poschmann & Hörschemeyer, 2010

Cantharidae Wedmann et al., 2010: 51 (and Fig. 2e)

FOSSILS & MINERALS

TYPE HORIZON. Late Oligocene (about 25 Ma: 24.79-24.56 or 28 Ma).

TYPE LOCALITY. Germany: Enspel Fossil-Lagerstätte.

PRESERVATION/LITHOLOGY. Compression/ impression. Crater lake, thick pyroclastics and intercalated sediments (black pelites).

COLLECTIONS. Coll. No. PE2007/5006-LS. Now in Generaldirektion Kulturelles Erbe RLP, Referat Erdgeschichte, Mainz (Germany) and will be deposited in the Naturhistorisches Museum Mainz/ Landessammlung für Naturkunde Rheinland-Pfalz (NHMM).

COMMENTS AND NOTES. One specimen with red pronotum and black elytra. Body length: about 17 mm?.

Cantharidae Kirejtshuk & Azar, 2013

Cantharidae Kirejtshuk & Azar, 2013a: 60; Kirejtshuk & Azar, 2013b: 103, 106, 110, 114

Hsiao et al. 2016: 120, 122; Peris et al. 2016a: 699, 700; Poinar & Fanti 2016: 1; Kirejtshuk & Ponomarenko 2009-2015.

TYPE HORIZON. Lower Cretaceous: Barremian - lowermost Aptian (about 125-135 Ma) but could be earlier.

TYPE LOCALITY. Central Lebanon: Hammana - Mdeirij outcrop.

PRESERVATION/LITHOLOGY. Inclusion in amber. Lebanese Amber.

COLLECTIONS. Temporarily deposited in the Muséum National d'Histoire Naturelle, Paris (MNHN), No. "1465", and No. "1012AB"

COMMENTS AND NOTES. One specimen with posterior body portions broken or missing (No. 1465) and one specimen (family *incertae sedis*) of a very small cantharoid larva (No. 1012AB).

Cantharidae Peris, Sánchez-García, Soriano & Delclòs, 2013

Cantharidae Peris et al., 2013: 74

Peris et al. 2016a: 699, Appendix 1; Rasnitsyn et al. 2016: Appendix A, Appendix B p. 7; Peris et al. 2016b: Poster (FIGURE 3. D, Table 1.).

TYPE HORIZON. Lower Cretaceous: Early-Middle Albian (about 110 Ma) or Upper Aptian - Lower Albian.

TYPE LOCALITY. Spain, Cantabria province, El Soplao.

PRESERVATION/LITHOLOGY. Inclusion in amber. Spanish Amber.

COLLECTIONS. El Soplao Cave Laboratory (Celis, Spain) Institutional amber collection No. CES-522.

COMMENTS AND NOTES. One specimen.

Illustrated in Peris et al. (2016b: poster: figure 3. D).

Cantharidae Katona, Kutasi, Papp & Tóth, 2014

Cantharidae Katona et al., 2014: 130, 138

TYPE HORIZON. Pliocene (4 Ma).

TYPE LOCALITY. Hungary, Veszprém County, near Pula (Pula alginite mine).

PRESERVATION/LITHOLOGY. Exoskeleton. Volcanic crater with alginite.

COLLECTIONS. Bakony Museum of the Hungarian Natural History Museum.

COMMENTS AND NOTES. The authors (Katona et al. 2014: 130, 138) say that is possible to find representatives of the family Cantharidae.

Cantharinae Bachofen-Echt, 1949

Cantharinae Bachofen-Echt, 1949: 108-109

COMMENTS AND NOTES. See under **Cantharidae** Bachofen-Echt, 1949.

Cantharinae Crowson, 1972

Cantharinae Crowson, 1972: 64

Spahr 1981b: 4; Ramsdale 2002: 205 [only as Cantharidae in Baltic amber].

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. Various specimens quite close to many recent genera.

Cantharinae Kubisz, 2001

Cantharinae Kubisz, 2001: 260

Ponar & Fanti: 1.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland ?: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of Amber Inclusions at the Department of Invertebrate Zoology, University of Gdańsk.

COMMENTS AND NOTES. One specimen.

Malthininae Kubisz, 2001

Malthininae Kubisz, 2001: 260

Ponar & Fanti: 1.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Poland ?: Baltic Sea coast.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Museum of Amber Inclusions at the Department of Invertebrate Zoology, University of Gdańsk.

COMMENTS AND NOTES. Two specimens.

Other data

-*Malthinus* sp. and *Malthodes* sp.: quite common in Baltic amber (Larsson 1978: 139).

-*Telephorus*?: three species from Oeningen, in whole or in part referable to *Telephorus* in Mr. Lacoe's collection (Scudder 1895: 120; Handlirsch 1906-1908: 740 [as *Cantharis* sp.]; Kirejtshuk & Ponomarenko 2009-2015 [Scudder 1895]).

-Indeterminated genera: four genera in Baltic amber (Klebs 1910: 237; Spahr 1981b: 15), but they could be Cantharidae, Melyridae or other families.

-*Téléphores*: various fossil species (Pictet 1854: 334).

-*Telephoridae*: Lias (Goss 1879: 17, 22, 36, reprint 1881: 130, 135, 149).

-*Telephoridae*: Lias (Goss 1880: 26, reprint 1881: 294).

-*Telephoridae*: 20 specimens in amber (Helm 1886: 271 [coll. Helm]; Spahr 1981b: 15).

-*Telephoriden*/*"Telephoridae"*: various specimens (Helm 1896: 228; Handlirsch 1906-1908: 741; Spahr 1981b: 15).

-*Telephoridae*: various specimens of Paleogene (62), Neogene (22) and Quaternary (3) (Handlirsch, 1920-1921: 288 (VII. Tabellarische / Tabelle I.); Kelner-Pillault 1970: 11 [Handlirsch]; Spahr 1981b: 15 [Kelner-Pillault]).

-*Cantharides*: cretaceous land (Oustalet 1874: 8).

-*Cantharidae*/*Telephoridae*: in Baltic amber (Klebs 1889: 53; Spahr 1981b: 15).

-*Canthariden*/*Cantharidae*: Tertiary: Oligocene and Miocene (Handlirsch 1906-1908: 1175, 1183 in Tabelle VII., 1277, 1288, 1291).

-*Cantharidae*/*Cantharididae*: in Baltic amber (Ander 1942: 25, 35, 36; Spahr 1981b: 15).

-*Cantharids*: few cantharids are known from the Danish collection (Larsson 1978: 140; Spahr 1981b: 4, 15).

-*Cantharidae*: cited by Brodie and Giebel (Goss 1879: 10, 27, reprint 1881: 123, 140).

-*Cantharidae*: in the Purbeck (Goss 1880: 26, reprint 1881: 294).

-*Cantharidae*: Scudder (1886: 69) cited this family in Meloidae but is considered (for me by mistake) as true Cantharidae by Spahr (1981b: 15).

-*Cantharidae*: Baltic amber (Katinas 1971: 33; Spahr 1981b: 15).

-*Cantharidae*: Baltic amber (Crowson 1972: 64; Spahr

1981b: 15).

-*Cantharidae*: one larva (Larsson 1978: 140; Spahr 1981b: 4) but is doubt if is a Cantharidae, a Lampyridae or other families.

-*Cantharidae*: Baltic amber (Willemstein 1987).

-*Cantharidae*: Eocene/Oligocene (Ponomarenko 1995: 168; Ramsdale 2002: 205 [Ponomarenko 1995]).

-*Cantharidae*: in Eocene Baltic amber (Grimaldi & Engel 2005: 386).

-*Cantharidae*: several species from Florissant (Nudds & Selden 2008: 222-223).

-*Cantharidae*: Bitterfeld amber (Dunlop 2010: 69).

-*Cantharidae*: Baltic amber (Weitschat & Wichard 2010: 113).

-*Cantharidae*: Rovno amber (Perkovsky et al. 2010: 136).

-*Cantharidae*: Baltic amber (Alekseev & Bukejs 2013: 14).

-*Cantharidae*: not present in Oise amber but numerous in Baltic amber (Kirejtshuk & Nel 2013: 178).

-*Cantharidae*: Caenozoic (Rasnitsyn et al. 2016: Appendix A, Appendix B p. 7).

-*Cantharididae*: 51 specimens in Baltic amber (Brues 1933: 395 Table I, 396 (Klebs collection); Spahr 1981b: 15).

-*Cantharididae*: six genera from the Tertiary fauna (Rohdendorf & Ponomarenko 1962: 258. English translation of 1991: 366).

-*Cantharoidea*: many species (Crowson et al. 1967: 527; Spahr 1981b: 4).

-*Malacodermes*: 65 amber specimens in Berend (sic!) collection at Danzig (Motschulsky 1857: 25; Spahr 1981b: 4).

-*Malacodermen*: of Oeningen and Baltisch amber (Assmann, 1870: 30-31; Spahr 1981b: 4).

-*Malacodermes*: era of Purbeck (Oustalet 1874: 25).

-*Malacodermata*: various specimens from Florissant (U.S.A.), not very well preserved (Scudder 1882: 292; Handlirsch 1906-1908: 741).

-*Malacodermatae* (sic!), *Telephoridae*, *Cantharidae* /: Baltic amber (Klebs list) housed at Königsberg Museum (Williamson 1932: 144; Spahr 1981b: 4, 15) / **soldier beetles**: Baltic amber (Williamson 1932: 142).

-*Elateridae* or *Telephoridae*: (Brodie 1875: 13. *Separatum*: not consulted).

-*Lampyridae*: three specimens in Lias of Schambelen - Switzerland (Scudder 1885: 796; Scudder 1886: 75 [Heer citation]; Scudder 1887: 796), and with the term Lampyridae, Scudder also was referring to Cantharidae.

-**Lampyridae**: twenty-one fossil species belonging to nine genera (Scudder 1900: 101), some of these are Cantharidae.

incertae sedis (probably not Cantharidae)

Genus †*Pseudotelephorus* Handlirsch, 1906: 454, Tafel XLI. Fig. 67. (caption)

†*Pseudotelephorus grandis* Cockerell, 1915

Pseudotelephorus grandis Cockerell, 1915: 479, 498 (explanation of plates), Plate 61. Fig. 3.

Champion et al. (eds.) 1916: 63; Handlirsch 1939: 71; Mitchell 2013 - EDNA The Fossil Insect Database [undetermined family]; The Paleobiology Database [undetermined family].

TYPE HORIZON. Triassic/Jurassic: Upper Rhaetian - Lower Hettangian (208-196.5 Ma).

TYPE LOCALITY. United Kingdom.

PRESERVATION/LITHOLOGY. Elytron(impression).

Marine shale/limestone.

COLLECTIONS. Collected by Brodie and purchased by British Museum (BMNH) in 1898. Now according Cockerell (1915: 469, 479) is deposited in the United States National Museum coll. Lacoce No. 3497 (Brodie leg.) now Cat. No. 61404 U.S.N.M.

COMMENTS AND NOTES. One elytron. Length: visible part about 11 mm (entirely probably about 12 mm.) x 4.0 mm. Base distinctly narrower than the middle. By Cockerell (1915: 478) this species is of the family Erotylidae. Handlirsch (1939: 71) cites this species under its new genus *Anancaeon* Handlirsch, which is a Coleoptera *incertae familiae* (e.g. Ponomarenko & Schultz 1988: 15) and that clearly is not a Cantharidae.

†*Pseudotelephorus haueri* (Giebel, 1856)

Telephorus Haueri Giebel, 1856a: 101 [under family Lampyridae], 400 (in the "Tabellarische")

Telephorus Haueri Giebel, 1856b: 177 [*nomen nudum*]

Brodie 1845: 101 (List), 124, Pl. 6. fig. 29 [unnamed]; Pictet 1854: 334 [unnamed]; Brodie 1873: 25 (tabular view) [*separatum*: 16 (tabular view)]; Goss 1879: 31, 32 (reprint 1881: 144, 145); Scudder 1885: 796 (fig. 1031) [under family Lampyridae]; Scudder 1886: 75-76 [under family Lampyridae]; Scudder 1887: 796 (fig. 1049) [under family Lampyridae]; Scudder 1891: 220 [under family Lampyridae]; Handlirsch 1906-1908: 454, Tafel XLI. Fig. 67. (and caption); Cockerell 1915: 478 [as *Pseudotelephorus haueiri*]; Handlirsch 1939: 71 [as *Pseudotelephorus Haueiri*]; Mitchell 2013 - EDNA The Fossil Insect Database [as *Telephorus haueri* and *Pseudotelephorus haueri*]; The Paleobiology Database [as *Pseudotelephorus haueri*].

TYPE HORIZON. Triassic: Rhaetian (208.5-201.3 Ma), sometimes considered Lower Lias.

TYPE LOCALITY. United Kingdom: Gloucestershire, Forthampton, Wainlode, Brown's Wood.

PRESERVATION/LITHOLOGY. Elytron (Mold/

impression). Marine horizon (Lilstock Formation). COLLECTIONS. Collected by Brodie and purchased by British Museum (BMNH) in 1898. Now according Cockerell (1915: 469, 478) is deposited in the United States National Museum, No. 3480 Lacoce coll. (Brodie leg.) now Cat. No. 61402 U.S.N.M.

COMMENTS AND NOTES. One elytron with very delicate punctures, redrawing in Handlirsch (1906-1908: Taf. XLI. Fig. 67.) Length: 8.0 mm (Handlirsch 1906-1908: 454) x 2.30 mm (The Paleobiology Database). By Cockerell (1915: 478) this species is not Cantharidae (Telephoridae) but Erotylidae for about 10 very delicate striae but not sharp, minute punctiform brown markings (not true punctures) that they may have been pigmented spots. Also for Handlirsch (1906-1908: 454) was an uncertain family and not *Telephorus*, in fact the same Handlirsch (1939: 71) cites this species under its new genus *Anancaeon* Handlirsch, which is a Coleoptera *incertae familiae* (e.g. Ponomarenko & Schultz 1988: 15) and that clearly is not a Cantharidae.

†*Pseudotelephorus punctulatus* Cockerell, 1915

Pseudotelephorus punctulatus Cockerell, 1915: 479, 498 (explanation of plates), Plate 61. Fig. 4.

Champion et al. (eds.) 1916: 63; Handlirsch 1939: 71; Mitchell 2013 - EDNA The Fossil Insect Database [undetermined family]; The Paleobiology Database [undetermined family].

TYPE HORIZON. Triassic/Jurassic: Upper Rhaetian - Lower Hettangian (208-196.5 Ma).

TYPE LOCALITY. United Kingdom.

PRESERVATION/LITHOLOGY. Elytron (impression). Marine shale/limestone.

COLLECTIONS. Collected by Brodie and purchased by British Museum (BMNH) in 1898. Now according Cockerell (1915: 469, 479) is deposited in the United States National Museum coll. Lacoce No. 3496 (Brodie leg.) now Cat. No. 61403 U.S.N.M.

COMMENTS AND NOTES. Portion of one elytron. Length: 7.5 mm (entirely probably little over 9 mm.) x 3.0 mm. Elytron with about 10 delicate striae and apex moderately acute, larger and broader than *P. haueri*, with the outer margin more convex (Cockerell 1915: 479). By Cockerell (1915: 478) this species is of the family Erotylidae. Handlirsch (1939: 71) cites this species under its new genus *Anancaeon* Handlirsch, which is a Coleoptera *incertae familiae* (e.g. Ponomarenko & Schultz 1988: 15) and that clearly is not a Cantharidae.

Genus †*Telephorium* Westwood, 1854: 395,

Plate XVII. Fig. 4.

†*Telephorium abgarus* Westwood, 1854

Telephorium Abgarus Westwood, 1854: 395 (explanation of Plate XVII.), Plate XVII. Fig. 4., 385-387 [family Telephoridae]

Giebel 1856a: 102 [under family Lampyridae]; Goss 1879: 27 (reprint 1881: 140); Scudder 1885: 796 [under family Lampyridae]; Scudder 1886: 76 [under family Lampyridae]; Scudder 1887: 796 [under family Lampyridae]; Scudder 1891: 220 [under family Lampyridae]; Handlirsch 1906-1908: 562, Tafel XLV. Fig. 71. (and caption); Coram & Jepson 2012: Figure 86 (p. 62), 64 [undetermined family]; Mitchell 2013 - EDNA The Fossil Insect Database [undetermined family]; The Paleobiology Database [undetermined family].

TYPE HORIZON. Early Cretaceous: Berriasian (145.0-139.8 Ma), Purbeck Group (Purbeck Formation).

TYPE LOCALITY. United Kingdom: Dorset, Purbeck, near Swanage (Durlston Bay).

PRESERVATION/LITHOLOGY. Elytron (compression). Fluvial-lacustrine limestone.

COLLECTIONS. Collected by Charles Willcox (probably in 1853; see The Paleobiology Database) and communicated by Brodie (Westwood 1854: 385, 395). Dorset County Museum (DORCM), Dorchester (UK), reposed in the British Museum of Natural History (BMNH).

COMMENTS AND NOTES. One elytron. Length: 7.00 mm (Handlirsch 1906-1908: 562) x 2.30 mm (The Paleobiology Database). Redrawn from the original illustration by Handlirsch (1906-1908: Tafel XLV. Fig. 71.) and by Coram & Jepson (2012: fig. 86). The elytron is equipped with some striae (10 by Handlirsch) that seems to indicate they do not belong to the family Cantharidae (Handlirsch 1906-1908: 562).

Cantharidae Sendel, 1742

Cantharidibus Sendel, 1742: 122-123, Tab. IV (Fig: 16-17) Fischer 1939: 95; Weitschat & Wichard 1998: 154, 155 (Tafel 57); Weitschat & Wichard: 2002: 152, 153 (Plate 57); Wichard & Greven 2009: Abb. 4 (285), 275 (replication of Sendel's table IV), 284-285 (also replication of Sendel's pages 122-123), 286

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Royal Museum in Dresden.

COMMENTS AND NOTES. They are the first known depictions of amber Cantharidae (and fossil Cantharidae) but at least one (Sendel 1742: Fig. 16) of these two Sendel's specimens is likely to be referred to family Cupedidae (Weitschat & Wichard 1998: 154, 155 (Tafel 57); 2002: 152, 153 (Plate

57); Wichard & Greven 2009: Abb. 4 (p. 285), 286). Personally, I suspect that the second specimen (Sendel 1742: Fig. 17) can not be a Cantharidae but probably Lampyridae or otherwise.

Cantharidae Berendt, 1845

Callidium Berendt, 1845: 56

Pictet 1854: 334; Giebel 1856a: 128; Zang 1905: 240; Handlirsch 1906-1908: 741; Larsson 1978: 139; Spahr 1981b: 15; Hieke & Pietrzeniuk 1984: 303, 305; Vitali 2009: 237 [see Vitali also for all citations as Cerambycidae].

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

TYPE LOCALITY. Probably Prussia.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COLLECTIONS. Berlin Museum (coll. Berendt).

COMMENTS AND NOTES. As stated in Vitali (2009: 237), one of the specimens of Berendt (1845: 56) called *Callidium* was transferred to families Cantharidae or Lampyridae by Zang (1905: 240), and agree with Vitali (2009: 237), probably, is the same specimen quoted by Hieke & Pietrzeniuk (1984: 305) as *Callidium*.

Cantharidae Menge, 1856

cantharidia Menge, 1856: 21

lampyrides Menge, 1856: 21

Larsson 1978: 139; Spahr 1981b: 15.

TYPE HORIZON. Eocene: Priabonian (38.0/37.2-33.9 Ma) sometimes considered Lutetian (48.6-40.4 Ma), or Early Oligocene (28.4-33.9 Ma).

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. 21 specimens of "cantharidia" and one of "lampyrides", whose assignment to Cantharidae is uncertain.

Cantharidae Bachofen-Echt, 1949

Junge Larve *Cantharidae* Bachofen-Echt, 1949: Abb. 104. (p. 115)

Larsson 1978: 150; Spahr 1981b: 15.

PRESERVATION/LITHOLOGY. Inclusion in amber. Baltic Amber.

COMMENTS AND NOTES. This specimen is was referred to family Cleridae by Larsson (1978: 150).

Cantharoidea Rasnitsyn & Ross, 2000

Superfamily *Cantharoidea* Rasnitsyn & Ross, 2000: 22

Grimaldi et al. 2002: 11 (Table 2); Hsiao et al. 2016: 122; Poinar & Fanti 2016: 1; Fanti & Ellenberger 2016: 166.

TYPE HORIZON. Early Cretaceous: Upper Albian (97-110 Ma).

TYPE LOCALITY. Myanmar.

PRESERVATION/LITHOLOGY. Inclusion in amber. Burmese Amber.

COLLECTIONS. Department of Palaeontology of The Natural History Museum, London, No. In.19123, and In.20169.

COMMENTS AND NOTES. One larva (In.19123) and one specimen doubt Cantharoidea (In.20169).

Cantharoid larva see under **Cantharidae** Kirejtshuk & Azar (2013b: 114).

Kirejtshuk & Ponomarenko 2009-2015.

Moreover these genera and/or species, although included the term “*cantharis*”, are for the habitus clearly belonging to other families, and not Cantharidae:

Cantharis lithophilus Wickham, 1914 = *Lytta lithophila* (Wickham, 1914) - Meloidae.

Melanocantharis Bode, 1953 - *incertae saeclis* Suborder Polyphaga.

Melanocantharis bicornuta Bode, 1953

Sphaerocantharis Bode, 1953 - *incertae saeclis* Suborder Polyphaga.

= *Sphaeracantharis* Bode, 1953: 368 (incorrect original spelling)

= *Mikrocarpides* Bode, 1953

= *Microcarpides* Bode, 1953: 369 (incorrect original spelling); Ponomarenko, 1992 (incorrect subsequent spelling. ICZN 1999 Art. 33.3)

= *Microcarpoides* Ponomarenko, 1990 (incorrect spelling)

= *Theornithion* Bode, 1953

Sphaerocantharis defossa Bode, 1953 (type species)

Sphaerocantharis striata (Bode, 1953) [*Cantharis* is feminine]

Theornithion striatum Bode, 1953

= *Mikrocarpides lineatus* Bode, 1953

Sphaerocantharis sibirica (Ponomarenko, 1990)

[*Cantharis* is feminine]

“*Microcarpoides*” *sibiricus* Ponomarenko, 1990

(see: Wickham 1914; Bode 1953; Ponomarenko 1990, 1992; Kirejtshuk et al. 2009-2015; The Paleobiology Database).

Viewing the illustration and the original descriptions, the species: *Miocaenia pectinicornis* Wickham, 1914 (from Chadronian lacustrine - large shale of Florissant, USA) and *Phausis fossilis* Beier, 1952 (from Chattian spring travertine of Halbopal at Luschitz [= Lužice], Czech Republic), belong very probably to the family Lampyridae (not Cantharidae), as in the descriptions. However, the genus *Phausis* LeConte (a long time in synonymy with *Lamprohiza* Motschulsky) is North American, and therefore the correct new combination is the following:

Lamprohiza fossilis (Beier, 1952) [n. comb.]

Phausis fossilis Beier, 1952: 131

4. Conclusion

The first representation, with doubts about the membership in the family Cantharidae, is present in Sendel (1742) and after we have only species to generic level or undeterminate (first were Burmeister 1832; Vollmar 1835; Burmeister 1836; Keferstein 1834; and particularly Hope 1836), with the first species surely Cantharidae, excluded *Cantharis* aff. *nigricans* Burmeister (1832) and cfr. *Autosilis nitidula* (Serres 1843), that have been described by Heer (1847). Now all the five subfamilies (Cantharinae, Chauliognathinae, Dysmorphocerinae, Malthininae, Silinae) and all the tribes (Cantharinae Cantharini, Cantharinae Podabrinii, Chauliognathinae Chauliognathini; Chauliognathinae; Ichthyurini; Malthininae Malchinini, Malthininae Malthinini, Malthininae Malthodini, Silinae Silini, Silinae Tytthonyxini, + the new tribe Malthininae Mimoplatycini) are known in the fossil record.

With this article I was able to survey 53 fossil species + *Cantharis* aff. *nigricans* and cfr. *Autosilis nitidula* + 6 species still living (*Cantharis figurata*, *Cantharis* cf. *paludosa*, *Cantharis rufa*, *Malthacus deceptus*, *Malthacus piniphilus* species group, *Podabrus* cfr. *alpinus*), + various specimens known to genus or family/subfamily level and particularly *Podistra* (*Absidia*) that does not currently has fossils at specific level.

I also provided several generic data, and the species included in Cantharidae but that probably are to be excluded of this family, as well as I provided a list of the fossil species in reference to the epoch and locality (Table 1.), and as can see the species of the family are known non-stop for every age from Cretaceous to Holocene. Particularly I furnished a list of subfossil species (Table 2.) that have been found to be 6 species (some cited in more locality) and other specimens at the generic or indeterminate level.

Considering that the first Elateriformia appeared on Permian (Toussaint et al. 2016) or at the end of Triassic / Lower Jurassic (Ponomarenko 1995; Chang et al. 2009b; McKenna et al. 2015) and the first Elateridae in the Early and Middle Jurassic (Chang et al. 2009a, 2009b), but for Toussaint et al. (2016) in Triassic as Elateroidea; is likely that the first arcaic representatives of the family Cantharidae have appeared and evolved in the Early Cretaceous (McKenna et al. 2015) or more probably, for me, in the Middle-Upper Giurassic; as in part demonstrated by the few indeterminate remains of this period of this family for the Lower Cretaceous of: Lebanese Amber (Kirejtshuk & Azar 2013a, 2013b: 135-125 Ma), Australia (Jell & Duncan 1986: 122.5-112 Ma) and Spanish Amber (Peris et al. 2016: about

FOSSILS & MINERALS

110 Ma) or by *Ornatomalthinus elvira*e (Poinar & Fanti 2016: 97-110 Ma), *Myamalycoerus vitalii* (Fanti & Ellenberger 2016: 97-110 Ma) and *Archaeomalthodes rosetta* (Hsiao et al. 2016: 97-110 Ma) from Early Cretaceous of Myanmar. After which, very probably, this family has suffered greatly the effects of the big mass extinction of the end of Cretaceous, with the likely extinction of many genera, and only in the first few million years of the Cainozoic were formed living genera, and in fact already in the Eocene (30-40 Ma) many current genera were already present (e.g. *Atalantycha*, *Cantharis*, *Podistra*, *Rhagonycha*, *Themus*, *Podabrus*, *Chauliognathus*, *Trypherus*, *Macrocerus*, *Malthinus*, *Malthodes*, *Polemius*, *Silis*).

Table 1. List of the fossil species in reference to epoch/period and locality

Cretaceous

Burmese Amber:

- Myamalycoerus vitalii* Fanti & Ellenberger, 2016 [97-110 Ma]
- Ornatomalthinus elvira*e Poinar & Fanti, 2016 [97-110 Ma]
- Archaeomalthodes rosetta* Hsiao, Ślipiński & Pang, 2016 [97-110 Ma]

Paleocene

Peng Chau Island, Lung Lok Shui (Hong Kong):

- Lithocantharis lunglokshuiensis* Lin, 1997 [66-56 Ma]

Peng Chau Island, Wong Ye Kok (Hong Kong):

- Wongyekokia angustris* Lin, 1997 [66-56 Ma]

Quebrada “El Griton”, “Sunchal Formation” (Argentina):

- Podabrus santaritensis* Cockerell, 1936 [65.5-56.0 Ma]

Eocene

Dominican Amber:

- Tytthonyx geiseri* Poinar & Fanti, 2016 [45-30 Ma to 20-15 Ma - Eocene or Miocene]

Baltic Amber:

- Cantharis aff. nigricans* Burmeister, 1832 [38/37.2-33.9 Ma]
- Cantharis (Cantharis) sucinonigra* Kuśka, 1992 [38/37.2-33.9 Ma]
- Cantharis (Cyrtomoptila) sucinokotejai* (Kuśka, 1996) [38/37.2-33.9 Ma]
- Electronycha prussica* Kazantsev, 2013 [38/37.2-33.9 Ma]
- Rhagonycha kryshtofovichi* (Yablokov-Khnzorian, 1960) [38/37.2-33.9 Ma]
- Rhagonycha sucinobaltica* Poinar & Fanti, 2016 [38/37.2-33.9 Ma]
- Sucinocanthalis baltica* Kuśka & Kania, 2010 [38/37.2-33.9 Ma]
- Sucinorhagonycha kulickae* Kuśka, 1996 [38/37.2-33.9 Ma]
- Themus (Haplothemus) pristinus* Kazantsev, 2013 [38/37.2-33.9 Ma]
- Cacomorphocerus cerambyx* Schaufuss, 1892 [38/37.2-33.9 Ma]
- Cacomorphocerus jantaricus* (Kuśka & Kania, 2010) [38/37.2-33.9 Ma]
- Macrocerus sucinopenninus* (Kuśka & Kania, 2010) [38/37.2-33.9 Ma]
- Malthinus danieli* Kuśka & Kania, 2010 [38/37.2-33.9 Ma]
- Malthodes ceranowiczae* Kuśka & Kupryjanowicz, 2005 [38/37.2-33.9 Ma]
- Malthodes kotejai* Kuśka & Kupryjanowicz, 2005 [38/37.2-33.9 Ma]
- Malthodes serafini* Kuśka & Kupryjanowicz, 2005 [38/37.2-33.9 Ma]
- Malthodes sucini* Kuśka & Kania, 2010 [38/37.2-33.9 Ma]
- Mimoplatycis notha* Kazantsev, 2013 [38/37.2-33.9 Ma] - also from Rovno Amber
- Curche pauli* Alekseev & Kazantsev, 2014 [38/37.2-33.9 Ma]

FOSSILS & MINERALS

-*Electrosilis minuta* Kazantsev, 2013 [38/37.2-33.9 Ma]

Rovno Amber:

-*Malthodes perkovskyi* Kazantsev, 2010 [38/37.2-33.9 Ma]

-*Malthodes rovnoensis* Kazantsev & Perkovsky, 2014 [38/37.2-33.9 Ma]

Florissant (USA):

-*Atalantycha humata* (Wickham, 1913) [37.2-33.9 Ma]

-*Rhagonycha hesperus* (Wickham, 1914) [37.2-33.9 Ma]

-*Podabrus cupesoides* Wickham, 1917 [37.2-33.9 Ma]

-*Podabrus florissantensis* Wickham, 1914 [37.2-33.9 Ma]

-*Podabrus fragmentatus* Wickham, 1914 [37.2-33.9 Ma]

-*Podabrus wheeleri* Wickham, 1909 [37.2-33.9 Ma]

-*Chauliognathus pristinus* Scudder, 1876 [37.2-33.9 Ma]

-*Trypherus aboriginalis* Wickham, 1913 [37.2-33.9 Ma]

-*Polemius crassicornis* Wickham, 1914 [37.2-33.9 Ma]

Aix-en-Provence (France):

-cfr. *Autosilis nitidula* (Fabricius) - Serres 1843 [? – Eocene/Oligocene] - extant species?

Oligocene

Brunstatt (France):

-*Malthodes obtusus* Förster, 1891 [33.9-28.4 Ma]

Rott (Germany):

-*Cantharis brodiei* (Heyden & Heyden, 1866) [28.4-23.0 Ma]

-*Cantharis caduca* (Heyden & Heyden, 1866) [28.4-23.0 Ma]

-*Cantharis carbonaria* (Heyden & Heyden, 1866) [28.4-23.0 Ma]

-*Cantharis exauctarata* (Heyden & Heyden, 1866) [28.4-23.0 Ma]

Miocene

Mexican (Chiapas) Amber:

-*Silis chiapasensis* Wittmer, 1963 [23.0-16.0 Ma]

Radoboj (Croatia):

-*Cantharis attavina* (Heer, 1847) [16-20 Ma]

Shandong, Shanwang, Linqu County (China):

-*Lycocerus guttula* (J. Zhang, 1989) [16-11.6 Ma]

-*Themus (Themus) capacis* (J. Zhang, 1989) [16-11.6 Ma]

-*Themus (Themus) thermophilus* (J. Zhang, 1989) [16-11.6 Ma]

-*Themus (Themus) trapezialis* (J. Zhang, Sun & X. Zhang, 1994) [16-11.6 Ma]

Oeningen (Germany):

-*Cantharis fragilis* (Heer, 1847) [10-12.7 Ma]

-*Cantharis macilenta* (Heer, 1865) [10-12.7 Ma]

-*Rhagonycha germari* (Heer, 1847) [10-12.7 Ma]

-*Rhagonycha tertiaria* (Heer, 1847) [10-12.7 Ma] - also from Radoboj

Pliocene

Lac Chambon (France):

-*Rhagonycha micans* Piton, 1939 [3.6-2.6 Ma]

Pleistocene

Kap København, North Greenland (Denmark):

-*Podabrus* cfr. *alpinus* (Paykull, 1798) - Böcher 1995 [1.8-2.2 Ma - Late Pliocene?] - extant species

Upton Warren, River Salwarpe (UK):

-*Cantharis figurata* Mannerheim - Coope et al. 1961 [about 42000 y/o] - extant species

-*Cantharis cf. paludosa* Fallén - Coope et al. 1961 [about 42000 y/o] - extant species

-*Cantharis rufa* Linnaeus - Coope et al. 1961 [about 42000 y/o] - extant species

Holocene

Northern Alaska, Ikpikpuk River (USA):

-*Malthacus deceptus* (Brown) - Matthews & Telka 1997 [about 9.670 y/o] - extant species

-*Malthacus piniphilus* species-group - Matthews & Telka 1997 [about 9.670 y/o] - extant species

Other localities that have returned Cantharidae not determined to specific level are:

Bitterfelder amber, Lebanese amber, Spanish amber, confluence of the River Limay with River Traful (Argentina), Puy-Saint-Jean (France), Randecker Maar (Germany), Niedersachsen - 3411 Willershausen clay pit (Germany), Enspel Fossil-Lagerstätte (Germany), Pula (Hungary), Titusville Site (USA), Rocky Mountain National Park - Roaring River (USA), Denali National Park and Preserve - Foraker River - Foraker Slump (USA), Bell Basin - Rock River exposure (Canada), Strathcona Fiord, Beaver Pond deposit (Canada), One Mile Creek (Canada), McAbee Fossil Beds (Canada), Koonwarra Fossil Bed (Australia), and probably also: Purbeck (UK) and Schambelen (Switzerland).

Table 2. Preliminary list of subfossil (Holocenic) species

Genus indet. Hall, 1990: 22 (Table 1), (5, 6)

Late Holocene (2000 ± 80 y/o) – USA: Maine, Starks, Sandy River [Sample 2] – Fragment, terrace sediments – One specimen.

Cantharidae Genus et sp. indet. Elias, Short & Waythomas, 1996: 298

[see in the fossils, precedent pages]

Gen. indet. Hellqvist & Eriksson, 2001: 149 (Table II.)

Holocene ($420 \pm 55 - 285 \pm 55$ y/o) – Sweden: Gamla Uppsala, Fyris River by Valsgärde – Insect, sediments/fluvial deposits – One specimen (Unit V2).

Cantharis obscura L. Mighall et al., 2002: 1179 (Table 5.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insect, sediments – One specimen (Sample 9).

Cantharis sp. Girling, 1979: 89 (Appendix 1.), on-line version: 29 (Appendix 1)

Holocene (5200-5100 y/o) – UK: Sweet Track, Drove site [sample T] – Insect, peat – One specimen.

Cantharis sp. Mighall et al., 2002: 1179 (Table 5.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insect, sediments – One specimen (Sample 13b).

Cantharis sp. Merritt et al., 2003: Appendix 1 Site 20 Table A1.13

Holocene (4120 ± 50 - 3855 ± 50 y/o. About 4000) – UK: Scotland, River Don valley east of Kintore, Nether Daugh site (NJ 800 160) – fragments, fine-grained sediments – Two heads, one right elytron.

Cantharis sp. Panova et al., 2003: 255 (Таблица 2.) [English translation: 227 (Table 2.)]

Holocene (9.230-4.000 y/o) – Russia: Polar Urals: eastern slope of the Rai-Iz massif ($66^{\circ}51'N - 65^{\circ}41'E$), that conventionally named Lake Pereval'noe, 260 m [Sample number 6] – Fragment, peat samples – One specimen.

Cantharis spp. Smith et al., 2005: 363 (Table 4)

Early Holocene (10.700-9300 y/o) for sample 5 and Middle/Late Holocene (3800-2900 to 2020-1530 y/o) for sample 40 – UK: Leicestershire, River Soar, Croft Quarry – Fragment, free organic peat – One specimen (Sample 5) and one specimen (Sample 40).

Cantharis sp. Kuzmina, 2015: 707 (Table 5.), 712

FOSSILS & MINERALS

Holocene (8400 y/o) – Russia: Sakha-Yacutia, Alazeya River – Insects, peat – Two specimens (Section 203, sample 5).

Rhagonycha fuscitibia Rey, 1891

R. femoralis (Brul.) Mighall et al., 2002: 1179 (Table 5.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insects, sediments – One specimen (Sample 9) and one specimen (Sample 19).

R. lignosa (Müll.) Mighall et al., 2002: 1179 (Table 5.), 1183 (Table 7.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insects, sediments – One specimen (Sample 13.1b) and one specimen (Sample 19).

Rhagonycha lignosa (Müll.) Smith et al., 2005: 363 (Table 4), 370

Middle/Late Holocene (3800-2900 to 2020-1530 y/o) – UK: Leicestershire, River Soar, Croft Quarry – Fragment, free organic peat – One specimen (Sample 40).

Rhagonycha lignosa (Müll.) Branch et al., 2012: 1152 (Table 3.), 1150

Holocene (6300-3900 y/o) – UK: Lower Thames Valley, Hornchurch Marshes – Insects, peat – One specimen (Sample Depth (m OD) -3.18 to -3.28).

Rhagonycha testacea (L.) Mighall et al., 2002: 1179 (Table 5.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insect, sediments – One specimen (Sample 13.1b).

Rhagonycha testacea (L.) Branch et al., 2012: 1152 (Table 3.), 1150

Holocene (6300-3900 y/o) – UK: Lower Thames Valley, Hornchurch Marshes – Insects, peat – Three specimens (Sample Depth (m OD) -3.18 to -3.28, -2.58 to -2.68, -2.28 to -2.38).

Podabrus alpinus (Payk.) Mighall et al., 2002: 1179 (Table 5.)

Holocene (2000-1700/1650 y/o) – UK: Wales, Cwmystwyth, Mine of Copa Hill – Insects, sediments – Three specimens (Sample 13.1b).

Podabrus sp. Elias et al., 1986: 137 (Table 1)

Scott et al. 2001: 201 (Appendix B).

Late Holocene (ca. 2400 y/o) – USA: Colorado, Rocky Mountain National Park, Roaring River – Fragments, Thick organic lens – specimen(s).

? *Podabrus* sp. Hall, 1990: 22 (Table 1), (5, 6)

Late Holocene (2000 ± 80 y/o) – USA: Maine, Starks, Sandy River [Sample 2] – Fragment, terrace sediments – One specimen.

Podabrus sp. Short et al., 1992: 386 (Table 2.)

Holocene (8580 ± 250 y/o) – USA: Alaska, Nushagak Lowland, Flounder Flat [Sample B] – Insect, peat – One specimen.

Silis ruficollis (F.) Girling, 1979: 89 (Appendix 1.), on-line version: 29 (Appendix 1)

Holocene (5200-5100 y/o) – UK: Sweet Track, Drove site [sample 21] – Insect, peat – One specimen.

Appendix 1. Etymology and type species of the fossil genera

Electronycha Kazantsev:

combination of Greek “*electron*” = amber + genus name *Rhagonycha*.

Gender: feminine.

Type species *Electronycha prussica* (Kazantsev 2013: 284), by original designation.

Lithocantharis Lin:

from Greek “*lithos*” = stone + *Cantharis* (lith- + canthar-).

Gender: feminine (not indicated in the description).

Type species *Lithocantharis lunglokshuiensis* (Lin 1997: 193), by original designation.

Myamalycoerus Fanti & Ellenberger:

combination of the specimen’s collection locality Myama and the extant genus *Lycocerus* Gorham

Gender: masculine.

Type species *Myamalycoerus vitalii* (Fanti & Ellenberger 2016: 166), by original designation.

Ornatomalthinus Poinar & Fanti:

from Latin “*ornatus*” = adorned (for sculpture elytra) and “*malthinus*” = soft/flabby.

Gender: masculine.

Type species *Ornatomalthinus elvira* (Poinar & Fanti 2016: 2), by original designation.

Sucinocantharis Kuśka & Kania:

from Latin “*sucinum*” = amber + *Cantharis*.

Gender: feminine.

Type species *Sucinocantharis baltica* (Kuśka & Kania 2010: 52), by original designation.

Sucinorhagonycha Kuśka:

combination from Latin “*sucinus*” sic! = amber + generic name *Rhagonycha*.

Gender: feminine

Type species *Sucinorhagonycha kulickae* (Kuśka 1996: 13), by original designation.

Curticantharis J. Zhang:

from Latin “*curti*” [from *curtus*] = brief/short + *Cantharis*.

Gender: feminine (not indicated in the description).

Type species *Curticantharis capacis* (J. Zhang 1989: 120, 422), by original designation.

Wongyekokia Lin:

from the locality Wong Ye Kok.

Gender: feminine (not indicated in the description).

Type species *Wongyekokia angustris* (Lin 1997: 194), by original designation.

Cacomorphocerus Schaufuss:

from Greek κακός = “malus” + μορφή = “forma” + κέρας = “cornu” (Schaufuss 1892).

Gender: masculine (not indicated in the description).

Type species *Cacomorphocerus cerambyx* (Schaufuss 1892), by monotypy.

Hoffeinsensia Kuśka & Kania:

in honor of Christel and Hans Werner Hoffeins from Hamburg, Germany.

Gender: feminine.

Type species *Hoffeinsensia jantarica* (Kuśka & Kania 2010: 50), by original designation.

Archaeomalthodes Hsiao, Ślipiński & Pang:

combination of the prefix *archaeo* = ancient, and the genus *Malthodes*.

Gender: masculine.

Type species *Archaeomalthodes rosetta* (Hsiao et al. 2016: 121), by original designation.

Mimoplatycis Kazantsev:

combination of Greek “*mimos*” = imitator + genus name *Platycis* (in reference to the pronotum).

Gender: feminine.

Type species *Mimoplatycis notha* (Kazantsev 2013: 288), by original designation.

Curche Alekseev & Kazantsev:

derived from Curche (Curcho, Kurche, Kurkas) = name of the god from the Old Prussian mythology. Gender: masculine.

Type species *Curche pauli* (Alekseev & Kazantsev 2014: 167), by original designation.

Electrosilis Kazantsev:

from Greek “*electron*” = amber + genus name *Silis*.

5. References

- ADAMONIS, S. (1994) - Entomofauna fosil de la Formacion Ventana (Eoceno) en el area de Confluencia, provincia de Neuquen. In: Resúmenes - VI Congreso Argentino de Paleontología y Bioestratigrafía, Trelew (1994): 48.
- ALEKSEEV, V. I. (2013) - The beetles (Insecta: Coleoptera) of Baltic amber: the checklist of described species and preliminary analysis of biodiversity. *Zoology and Ecology*, 23(1): 5-12.
- ALEKSEEV, V. I. & BUKEJS, A. (2013) - The beetles (Insecta: Coleoptera) of Baltic amber: the present state of systematic knowledge (p. 14). 7th International Conference "Research and conservation of biological diversity in Baltic region", Book of Abstracts, Daugavpils, 25-27 April, 2013.
- ALEKSEEV, V. I. & KAZANTSEV, S. V. (2014) - New fossil soldier beetle (Coleoptera: Cantharidae) from Baltic amber. *Baltic Journal of Coleopterology*, 14(2): 167-170.
- ANDER, K. E. V. (1942) - Die Insektenfauna des Baltischen Bernsteins nebst damit verknüpften zoogeographischen Problemen. *Lunds Universitets Årsskrift, N. F. Avd. 2, 38(4)* (Kunglig Fysiografiska Sällskapets Handlingar, N. F., 53(4): 1-82, 1 l. incl. illus. (maps) tables.
- ARILLO, A. (2007) - Paleoethology: fossilized behaviours in amber. *Geologica Acta*, 5(2): 159-166.
- [AUTHORLESS] - Inventaire de la faune fossile identifiée dans la collection d'ambre de Chiapas de Berkeley (Californie) et qui correspond au remarquable travail d'identification de Paul Hurd (1962)... *Le Mag - AMP*, 3: 38-42.
- ASSMANN, A. (1870) - Beiträge zur Insekten-Fauna der Vorwelt. *Zeitschrift für Entomologie, Vereins für Schlesische Insektenkunde zu Breslau*, 2(1): 3-62, 1 pl.
- BACHOFEN-ECHT, A. F. (1949) - Der Bernstein und seine Einschlüsse. Springer-Verlag, Wien, 204 pp.
- BEIER, M. W. P. (1952) - Miozäne und oligozäne Insekten aus Österreich und den unmittelbar angrenzenden Gebieten. *Sitzungsberichte der Akademie der Wissenschaften mathematisch-naturwissenschaftliche Klasse*, 161: 129-134.
- BERENDT, G. K. (1845) - Die organischen Bernstein-Einschlüsse im Allgemeinen. (pp. 41-60). In: GOEPPERT, H. R. & BERENDT, G. K.: Der Bernstein und die in ihm befindlichen Pflanzenreste der Vorwelt. In: BERENDT, G. K. (Hrsg.): Organische Reste im Bernstein. Nicolai, Berlin.
- BINDER, H. (2008) - Bernstein – Einblicke in vergangene Ökologie II: Käfer beim Einsatz chemischer Waffen ertappt. *Studium Integrale Journal*, 15 (1): 47.
- BÖCHER, J. (1995) - Palaeoentomology of the Kap København Formation, a Plio-Pleistocene sequence in Peary Land, North Greenland. *Meddelelser om Grønland, Geoscience*, 33: 82 pp.
- BODE, A. (1953) - Die Insektenfauna des ostniedersächsischen oberen Lias. *Palaeontographica, Abt. A*, 103: 1-375, 1 Karte, 15 t.
- BOUCOT, A. J. & POINAR JR., G. O. (2010) - Fossil Behavior Compendium. CRC Press, Boca Raton, 424 pp.
- BRANCH, N. P., BATCHELOR, C. R., CAMERON, N. G., RUSSELL COOPE, G., DENSEM, R., GALE, R., GREEN, C. P. and WILLIAMS, A. N. (2012) - Holocene environmental changes in the Lower Thames Valley, London, UK: Implications for understanding the history of Taxus woodland. - *The Holocene*, 22(10): 1143-1158.
- BRODIE, P. B. (1845) - A history of the fossil insects in the secondary rocks of England. Accompanied by a particular account of the strata in which they occur, and of the circumstances connected with their preservation. John Van Voorst, London, [xviii], errata, 1-130 + 11 plates + 4 pp.
- BRODIE, P. B. (1873) - The Distribution and Correlation of Fossil Insects, and the supposed occurrence of Lepidoptera and Arachnidae in British and Foreign Strata, chiefly in the Secondary rocks. Annual Report of the Warwickshire Natural History and Archaeological Society, 37: 12-28. [separatum: read at the Annual Meeting of the Warwickshire Natural History and Archaeological Society, Held at the Museum. Warwick, April 18th, 1873, 19 pp.].
- BRODIE, P. B. (1875) - On the Lower Lias at Evington and Kineton, and on the Rhaetian in that neighbourhood and their further extension in Leicestershire, Nottinghamshire, Lincolnshire, Yorkshire, and Cumberland. Annual Report of the Warwickshire Natural History and Archaeological Society, 39: 6-17. [separatum: (not consulted) fide Scudder, 1890: read at the Annual Meeting of the Warwickshire Natural History and Archaeological Society, held at the museum. Warwick, April 2nd, 1875, 14 pp.].

- BROWN, W. J. (1940) - Some new species of Cantharidae and Chrysomelidae (Coleoptera). *The Canadian Entomologist*, 72: 161-166.
- BRUCE ARCHIBALD, S., BOSSERT, W. H., GREENWOOD, D. R., and FARRELL, B. D. (2010) - Seasonality, the latitudinal gradient of diversity, and Eocene insects. *Paleobiology*, 36(3): 374-398 + Online Supplementary Material, 86 pp.
- BRUES, C. T. (1933) - Progressive change in the insect population of forests since the Early Tertiary. *The American Naturalist*, 67 (No. 712): 385-406.
- BURMEISTER, K. H. K. (1832) - *Handbuch der Entomologie*. Vol. 1. G. Reimer, Berlin, xvi + 696 pp. + Tables: 22 pp., 16 t.
- BURMEISTER, K. H. K. (1836) - Insects of a former world (pp. 574-581). A Manual of Entomology, Edward Churton Public Library, London, 1 p., xii + 654 pp. [translated from the German of Dr. Hermann Burmeister by W. E. Shuckard].
- CARPENTER, F. M. (1992) - Treatise on Invertebrate Paleontology. Part R. Arthropoda 4. Volume 4: Superclass Hexapoda. Boulder: The Geological Society of America, pp. i-ii, 279-655 [1992.12.09].
- CARUS, J. V. (1892) ed. - Litteratur. *Zoologischer Anzeiger*, 15: viii + 336 pp.
- CHAMPION, G. C., COLLIN, J. E., FOWLER, W. W., LLOYD, R. W., PORRITT, G. T. & WALKER, J. J. (1916) eds. - British Fossil insects. *The Entomologist's Monthly Magazine*, 52: 63-64.
- CHANG, H., KIREJTSUK, A. G., REN, D., and SHIH, C. (2009a) - First fossil click beetles from the Middle Jurassic of Inner Mongolia, China (Coleoptera: Elateridae). *Annales Zoologici (Warszawa)*, 59(1): 7-14.
- CHANG, H., ZHAO, Y., REN, D. (2009b) - New fossil elaterids (Insect: Coleoptera: Polyphaga: Elateridae) from the Middle Jurassic of Inner Mongolia, China. *Progress in Natural Science*, 19: 1433-1437.
- CIFUENTES-RUIZ, P., VEGA, F. J., CEVALLOS-FERRIZ, S. R. S., GONZÁLEZ-SORIANO, E., ZARAGOZA-CABALLERO, S. and GARIBAY-ROMERO, L. (2007) - Oligocene scorpion and insects (Plecoptera and Coleoptera) from the Los Ahuehuetes Locality, Puebla, Mexico. *Ameghiniana*, 44(4): 673-679.
- COCKERELL, T. D. A. (1915) - British fossil insects. *Proceedings of the United States National Museum*, 49(No. 2119) [1916]: 469-499, 6 plates (pls. 60-65). [this article appears in 1916 in the Proceedings, but the separatum of 30 pp. and plates, with same pagination, is output earlier: December 11, 1915 as British fossil insects, Washington, Government Printing Office].
- COCKERELL, T. D. A. (1936) - The fauna of the Sunchal (or Margas Verdes) Formation, Northern Argentina. *American Museum Novitates*, 886: 1-9.
- CONG, S., ASHWORTH, A. C., SCHWERT, D. P., and TOTTEN, S. M. (1996) - Fossil Beetle Evidence for a Short Warm Interval near 40,000 yr B.P. at Titusville, Pennsylvania. *Quaternary Research*, 45: 216-225.
- COOPE, G. R., SHOTTON, F. W., and STRACHAN, I. (1961) - A Late Pleistocene fauna and flora from Upton Warren, Worcestershire. With an appendix on the genus Pisidium at Upton Warren by S. P. DANCE. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 244: 379-421 + plates 19-22.
- CORAM, R. A. & JEPSON, J. E. (2012) - Fossil Insects of the Purbeck Limestone Group of Southern England: Palaeoentomology from the dawn of the Cretaceous. Siri Scientific Press, Manchester, 144 pp.
- COSSMANN, A. É. M. (1918) - New Species of fossil Beetles from Florissant, Colorado, by H. F. Vickham. *Revue critique de Paléozoologie*, 22(1-2): 13-14.
- CROWSON, R. A. (1972) - A review of the classification of Cantharoidea (Coleoptera), with the definition of two new families, Cneoglossidae and Omethidae. *Revista de la Universidad de Madrid*, 21: 35-77.
- CROWSON, R. A., ROLFE, W. D. I., SMART, J., WATERSTON, C. D., WILLEY, E. C., WOOTTON, R. J. (1967) - Arthropoda: Chelicera, Pycnogonida, Palaeoisopus, Myriapoda and Insecta. *The Fossil Record*, Pt. II, Chapter 19: 499-534. London (Geological Society).
- DELKESKAMP, K. (1939) - *Coleopterorum Catalogus*. Pars 165: Cantharidae. - S. Schenkling (ed.), Dr. W. Junk, 's-Gravenhage, 357 pp.
- DELKESKAMP, K. (1977) - *Coleopterorum Catalogus Supplementa*. Pars 165, Fasc. 1: Cantharidae. - J. A. Wilcox (ed.), Dr. W. Junk bv. Publishers, The Hague, 485 pp.
- DOUGLAS, S. D., and STOCKEY, R. A. (1996) - Insect fossils in middle Eocene deposits from British Columbia and Washington State: faunal diversity and geological range extensions. *Canadian Journal of Zoology*, 74(6): 1440-1157.

- DUNLOP, J. A. (2010) - Bitterfeld Amber (pp. 58-69). In: PENNEY, D. (Ed.). Biodiversity of fossils in Amber from the major world deposits. Siri Scientific Press, Manchester, 304 pp.
- ELIAS, S. A., SHORT, S. K. and CLARK, P. U. (1986) - Paleoenvironmental interpretations of the Late Holocene, Rocky Mountain National Park, Colorado, U.S.A. *Revue de Paléobiologie*, 5(1): 127-142.
- ELIAS, S. A., SHORT, S. K., and WAYTHOMAS, C. F. (1996) - Late Quaternary Environments, Denali National Park and Preserve, Alaska. *Arctic*, 49(3): 292-305.
- ENGEL, M. S. (2004) - Arthropods in Mexican amber (pp. 175-186). In: LLORENTE-BOUSQUETS, J., MORRONE, J. J., YÁÑEZ-ORDÓÑEZ, O., VARGAS-FERNÁNDEZ, I. (eds.). *Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia una Síntesis de su Conocimiento* (Volume IV). Universidad Nacional Autónoma de México, México DF, México, 790 pp.
- ETH ZÜRICH - Insekten aus der Fundstelle Öhningen (Miozän) in der Sammlung O. Heer des Departements Erdwissenschaften der ETH Zürich. ETH Zürich, Zürich, 20 pp. (unpaginated).
- FANTI, F. (2014) - Catalogo critico delle Cantharidae d'Italia (Insecta, Coleoptera). *Memorie della Società Entomologica Italiana*, 91(1-2): 61-132.
- FANTI, F. & ELLENBERGER, S. (2016) - Myamalyccerus vitalii: A new genus and species of soldier beetle in Burmese amber (Coleoptera Cantharidae). *Cretaceous Research*, 71 (2017): 166-169. dx.doi.org/ 10.1016/j.cretres.2016.11.019 [Available online 5 December 2016].
- FISCHER, C. W. (1939) - Mineralogie in Sachsen von Agricola bis Werner. Die ältere Geschichte des Staatlichen Museums für Mineralogie und Geologie zu Dresden (1560-1820). C. Heinrich, Dresden, viii + 347 pp., (24 tables).
- FÖRSTER, B. (1891) - Die Insekten des "Plattigen Steinmergels" von Brunstatt. Abhandlungen zur Geologischen Specialkarte von Elsass-Lothringen, 3(5): 335-593 + [1], 6 t. (+ 6), + [1].
- GERSDORF, E. (1971) - Weitere Käfer (COLEOPTERA) aus dem Jungtertiär Norddeutschlands. [2. Teil]. *Geologisches Jahrbuch*, 88: 629-669.
- GIEBEL, C. G. A. (1852) - Deutschlands Petrefacten. Ein systematisches Verzeichniss aller in Deutschland und den angrenzenden Ländern vorkommenden Petrefacten nebst Angabe der Synonymen und Fundorte. Verlag von Ambrosius Abel, Leipzig, xiii + 706 pp.
- GIEBEL, C. G. A. (1856a) - Fauna der Vorwelt mit steter Berücksichtigung der lebenden Thiere. Zweiter Band: Gliederthiere. Erste Abtheilung: Insecten und Spinnen. F. A. Brockhaus, Leipzig, xviii + 511 pp.
- GIEBEL, C. G. A. (1856b) - Geologische Uebersicht der vorweltlichen Insecten. *Zeitschrift für die Gesammten Naturwissenschaften*, 8: 174-188.
- GIRLING, M. A. (1979) - Fossil Insects from the Sweet Track. *Somerset Levels Papers*, 5: 84-93 [on-line version: 36 pp.]
- GOSS, H. (1878) - Three Papers on Fossil Insects, and the British and Foreign Formations in which Insect Remains have been detected. No. 1. The Insect Fauna of the Recent and Tertiary Periods, and the British and Foreign Formations of those Periods in which Insect Remains have been detected. London, 65 pp. [reprint: GOSS, H. (1878) in: *Proceedings of the Geologists' Association*, 5(6): 282-343] GOSS, H. (1879) - Three Papers on Fossil Insects, and the British and Foreign Formations in which Insect Remains have been detected. No. 2. The Insect Fauna of the Secondary or Mesozoic Period, and the British and Foreign Formations of that Period in which Insect Remains have been detected. London, 37 pp. [reprint: GOSS, H (1881) in: *Proceedings of the Geologists' Association*, 6(3): 116-150].
- GOSS, H. (1880) - Three Papers on Fossil Insects, and the British and Foreign Formations in which Insect Remains have been detected. No. 3. The Insect Fauna of the Primary or Palaeozoic Period, and the British and Foreign Formations of that Period in which Insect Remains have been detected. London, 32 pp. [reprint: GOSS, H (1881) in: *Proceedings of the Geologists' Association*, 6(6): 271-300].
- GRABAU, A. W. & SHIMER, H. W. (1910) - North American index fossils. Vol. II. A. G. Seiler & Company, New York, xiv + 1 [errata] + 909 pp.
- GRIMALDI, D. A. & ENGEL, M. S. (2005) - Evolution of the Insects. Cambridge University Press, Cambridge, UK, xv + 755 pp.
- GRIMALDI, D. A., ENGEL, M. S., NASCIMBENE, P. C. (2002) - Fossiliferous Cretaceous Amber from Myanmar (Burma): Its Rediscovery, Biotic Diversity, and Paleontological Significance. *American Museum Novitates*, 3361, 71 pp.
- HALL, H. A. (1990) - Paleoenvironmental analysis of a late-holocene subfossil Coleopteran fauna from Starks,

Maine. Senior Scholar Papers. Paper 110, 24 pp.

HANDLIRSCH, A. P. J. (1906-1908) - Die Fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen. Leipzig, Verlag: Wilhelm Engelmann, 1 p. + ix + 1430 pp. illustrated, diagrams (part fold) + atlas. [issued in parts between 1906 and 1908. Probably: pp. 1-640 in 1906, 641-960 in 1907, 961-1430 in 1908].

HANDLIRSCH, A. P. J. (1920-1921) - Siebentes Kapitel. Palaeontology. In: SCHRÖDER, C. (Ed.), Handbuch der Entomologie. 3. Band. Gustav Fischer, Jena, pp. 117-306. [issued in 1920 (pp. 113-208), 1921 (pp. 209-306)].

HANDLIRSCH, A. P. J. (1939) - Neue Untersuchungen über die fossilen Insekten mit Ergänzungen und Nachträgen sowie Ausblicken auf phylogenetische, palaeogeographische und allgemein biologische Probleme. II Teil. Annalen des Naturhistorischen Museums in Wien, 49 [1938]: 1-240, 16 pls.

HEER, O. (1847) - Die Insektenfauna der Tertiärgebilde von Oeningen und von Radoboj in Croatia. Erste Abtheilung: Käfer. Erster Theil: Käfer. Verlag von Wilhelm Engelmann, Leipzig, 229 pp. + [1] + 8 pls. [reprinted in: Neue Denkschriften der Allgemeinen Schweizerischen Gesellschaft für die gesammten Naturwissenschaften. Nouveaux Mémoires de la Société Helvétique des Sciences Naturelles, 8].

HEER, O. (1865) - Die Urwelt der Schweiz. Verlag von Friedrich Schulthess, Zürich, xxx + 622 pp., 11 pls. [French translation: HEER, O. (1872). English translation: HEER, O. (1876). German reprint: HEER, O. (1883)].

HEER, O. (1872) - Le Monde primitif de la Suisse. H. Georg, Libraire-Editeur, Genève et Bâle, frontispice + xvi + 801 pp. (i.e. 781) + xi (i.e. 20) t. (2 plié), cartes. [French translation of Heer, 1865 by Isaac Demole].

HEER, O. (1876) - The primaeval world of Switzerland. Vol. II. James Heywood (Ed.), London, Longmans, Green, and Co., 346 pp. [English translation of Heer, 1865].

HEER, O. (1883) - Die Urwelt der Schweiz. Zweiten, umgearbeiteten und vermehrten Auflage. Verlag von Friedrich Schulthess, Zürich, xix + 713 pp. + 1 + xii t. + 1 karte. [reprint of Heer, 1865].

HELLQVIST, M. & ERIKSSON, J. A. (2001) - Land-use history in Gamla Uppsala, southeast Sweden, established on subfossil Coleoptera remains and pollen in fluvial deposits. Grana, 40: 142-153.

HELM, O. (1886) - Mittheilungen über Bernstein. XIII. Ueber die Insecten des Bernsteins. Schriften der Naturforschenden Gesellschaft in Danzig, N. F., 6(3): 267-277.

HELM, O. (1896) - Beiträge zur Kenntniss der Insecten des Bernsteins. Schriften der Naturforschenden Gesellschaft in Danzig, N. F., 9(1): 220-231.

HELM, O. (1897) - Thierische Einschlüsse im Succinit. Schriften der Naturforschenden Gesellschaft in Danzig, N. F., 9(2): 88-89.

HENWOOD, A. A. (1992) - Soft-part preservation of beetles in Tertiary amber from the Dominican Republic. Palaeontology, 35(4): 901-912.

HEYDEN, C. H. G. von & HEYDEN, L. F. J. D. von (1866) - Käfer und Polypen aus der Braunkohle des Siebengebirges. Palaeontographica, 15: 131-156 (+ p. 157: Dipteren-Larve aus dem Tertiär-Thon von Nieder-Flörsheim in Rhein-Hessen), t. 22-24. [separatum under the title: Käfer und Polypen aus der Braunkohle des Siebengebirges. – Dipteren-Larve aus dem Tertiär-Thon von Nieder-Flörsheim in Rhein-Hessen. Besonderer Abdruck aus den Palaeontographicis, XV. Verlag von Theodor Fischer, Cassel, 1866, pp. 1-29, t. 1-3].

HEYDEN, L. F. J. D. von (1870) - Fossile Dipteren aus der Braunkohle von Rott im Siebengebirge. Palaeontographica, 17: 237-266 (p. 265-266: Nachträge), t. 44-45. [separatum: Cassel, 1870, pp. 1-30 (p. 29-30: Nachträge), t. 1-2].

HEYNDERYCX, J. (2004) - Insecten uit het Baltisch amber. Atalanta VZW, 30(1): 2-9. [in Dutch. Author's name of article on page 1].

HIEKE, F. & PIETRZENIUK, E. (1984) - Die Bernstein - Käfer des Museums für Naturkunde, Berlin (Insecta, Coleoptera). The Amber Beetles (Insecta, Coleoptera) of the Museum of Natural History, Berlin. Mitteilungen aus dem Zoologischen Museum in Berlin, 60(2): 297-326.

HOFFEINS, C. (2008) - Inkluzje zwierzęce i roślinne w kolekcji Otto Helma (1826-1902), Gdańsk. Animal and plant inclusions in the collection of Otto Helm (1826-1902), Gdańsk. Bursztynisko, 31, Special Edition, Gdańsk, Gdynia, 5-6.XII.2008: 36-44. [bilingual: in Polish and English].

HOPE, F. W. (1836) - Observations on Succinic Insects. The Transactions of the Entomological Society of London, 1: 133-147.

HSIAO, Y. and OKUSHIMA, Y. (2015) - A New Species of the Genus *Lycocerus* (Coleoptera, Cantharidae)

- from Taiwan. Elytra, Tokyo, New Series, 5(1): 109-113.
- HSIAO, Y., ŚLIPIŃSKI, A., DENG, C., PANG, H. (2016) - A new genus and species of soldier beetle from Upper Cretaceous Burmese amber (Coleoptera, Cantharidae, Malthininae). *Cretaceous Research*, 69 (2017): 119-123. [doi: 10.1016/j.cretres.2016.09.002 (Available online 15 September 2016)].
- HURD JR., P. D., SMITH, R. F., DURHAM, J. W. (1962) - The fossiliferous amber of Chiapas, Mexico. *Ciencia (Mexico)*, 21(3): 107-118 + 2 plates.
- INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE (1916a) - K Palaeontology. Royal Society of London, London, viii + 146 pp.
- INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE (1916b) - N Zoology. Royal Society of London, London, 342 pp.
- INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE (1919) - K Palaeontology. Royal Society of London, London, viii + 130 pp.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE - ICBN (1999) - International Code of Zoological Nomenclature, Fourth Edition. The International Trust for Zoological Nomenclature c/o The Natural History Museum, London, 306 pp.
- JANZEN, J.-W. (2002) - Arthropods in Baltic Amber. Ampyx-Verlag, 167 pp.
- JELL, P. A. (2004) - The fossil insects of Australia. *Memoirs of the Queensland Museum*, 50(1): 1-124.
- JELL, P. A. and DUNCAN, P. M. (1986) - Invertebrates, mainly insects, from the freshwater, Lower Cretaceous, Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association of Australasian Palaeontologists*, 3: 111-205.
- KANIA, I. (2015) - Autoreferat. Załącznik nr 2. Rzeszów (2015), 33 pp. [in Polish].
- KATINAS, V. I. (1971) - Amber and amber-bearing deposits of the southern Baltic area. Publishing Office "Mintis", Transactions, Volume 20, Vilnius, 156 pp. [in Russian].
- KATONA, L. T., KUTASI, C., PAPP, B. & TÓTH, S. (2014) - Újabb szenzációs őslénytani leletek a pulai alginítbányából. Further remarkable palaeontological finds at the alginite quarry in Pula. *Annales Historico-Naturales Musei Nationalis Hungarici*, 106: 117-140. [in Hungarian and English].
- KAZANTSEV, S. V. (2010) - New Malthodes (Insecta: Cantharidae: Coleoptera) from the Rovno Amber (Upper Eocene of Ukraine). *Russian Entomological Journal*, 19(2): 105-107.
- KAZANTSEV, S. V. (2013) - New taxa of Baltic amber soldier beetles (Insecta: Coleoptera: Cantharidae) with synonymic and taxonomic notes. *Russian Entomological Journal*, 22(4): 283-291.
- KAZANTSEV, S. V. & BRANCUCCI, M. (2007) - Family Cantharidae Imhoff, 1856 (1815). Pp. 234-298. In: LÖBL, I. & SMETANA, A. (eds.). Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea. Stenstrup: Apollo Books, 935 pp.
- KAZANTSEV, S. V. & PERKOVSKY, E. E. (2014) - A new Malthodes and some other interesting soldier beetles (Coleoptera: Cantharidae) from Late Eocene Rovno amber. *Russian Entomological Journal*, 23(2): 113-116.
- KEFERSTEIN, C. (1834) - Die Naturgeschichte des Erdkörpers in ihren ersten Grundzügen dargestellt. Vol. 2. Die Geologie und Paläontologie. Friedrich Fleischer, Leipzig, 896 pp.
- KELNER-PILLAULT, S. (1970) - L'ambre balte et sa faune entomologique avec description de deux Apoides nouveaux. *Annales de la Société entomologique de France* (N. S.), 6(1): 3-24.
- KIREJTSCHUK, A. G. & AZAR, D. (2013a) - Results of study of Coleoptera (Insecta) from the Lower Cretaceous Lebanese amber (p. 60). The 6th International Congress on Fossil Insects, Arthropods and Amber, Abstract Book, Byblos, Lebanon, April 14-18, 2013 [Oral].
- KIREJTSCHUK, A. G. & AZAR, D. (2013b) - Current knowledge of Coleoptera (Insecta) from the Lower Cretaceous Lebanese amber and taxonomical notes for some Mesozoic groups. *Terrestrial Arthropod Reviews*, 6: 103-134.
- KIREJTSCHUK, A. G. & NEL, A. (2013) - Current knowledge of Coleoptera (Insecta) from the Lowermost Eocene Oise amber. *Insect Systematics & Evolution*, 44: 175-201.
- KIREJTSCHUK, A. G. & PONOMARENKO, A. G. (2009-2015) - Taxonomical list of fossil beetles of the suborder Scarabeina (part2). <https://www.zin.ru/Animalia/Coleoptera/eng/paleosy1.htm> [last visit: December 2016].
- KIREJTSCHUK, A. G., PONOMARENKO, A. G., ZHERIKHIN, V. V. (2009-2015) - Taxonomic list of fossil beetles suborder Scarabeina (part4). <https://www.zin.ru/Animalia/Coleoptera/eng/paleosy3.htm> [last visit:

December 2016].

- KLAUSNITZER, B. (2003) - Buchbesprechungen. Entomologische Nachrichten und Berichte, 47: 54-55.
- KLEBS, E. H. R. (1889) - Aufstellung und Katalog des Bernstein-Museums von Stantien & Becker, Königsberg i. Pr. Nebst einer kurzen Geschichte des Bernsteins. Hartung, Königsberg, 103 pp.
- KLEBS, E. H. R. (1910) - Über Bernsteineinschlüsse im allgemeinen und die Coleopteren meiner Bernstein-sammlung. Schriften der Physikalisch-ökonomischen Gesellschaft zu Königsberg in Pr., 51: 217-242.
- KORSCHESKY, R. (1939) - Abbildungen und Bemerkungen zu vier Schaufuß'schen Coleopteren aus dem deutschen Bernstein. Arbeiten über morphologische und taxonomische Entomologie, 6 (1): 11-12 + t. 1.
- KOSMOWSKA-CERANOWICZ, B., KWIATKOWSKA, K., PIELIŃSKA, A. (2001) - The amber collection of the Museum of the Earth, Polish Academy of Sciences, as a source of multidisciplinary research (in commemoration of the collection's 50th anniversary). In: BUTRIMAS, A. (ed.). Baltic amber: Proceedings of the International Interdisciplinary Conference Baltic Amber in Natural Sciences, Archaeology and Applied Arts. Vilniaus Daile's akademijos leidykla, Vilnius (Acta Academiae Artium Vilnensis), 22
- KUBISZ, D. (2000) - Fossil beetles (Coleoptera) from Baltic amber in the collection of the Museum of Natural History of ISEA in Kraków. Polskie Pismo Entomologiczne, 69: 225-230.
- KUBISZ, D. (2001) - Beetles in the collection of the Museum of Amber Inclusions, University of Gdańsk, with description of *Colotes sambicus* sp. n. (Coleoptera: Melyridae). Polskie Pismo Entomologiczne, 70: 259-265.
- KUŚKA, A. (1992) - Three new species of beetles (Coleoptera: Cantharidae, Anobiidae, Curculionidae) from the Baltic amber. Annals of the Upper Silesian Museum - Entomology, 3: 107-113.
- KUŚKA, A. (1994) - Baltic amber inclusions of Lycidae and Cantharidae from the collection of the Museum of the Earth. Inclusion Wrostek, 14: 3, 6.
- KUŚKA, A. (1996a) - New coleopterans (Cantharidae, Curculionidae) from the collection of the Museum of the Earth. Inclusion Wrostek, 22: 19.
- KUŚKA, A. (1996b) - New beetle species (Coleoptera: Cantharidae, Curculionidae) from the Baltic amber. Prace Muzeum Ziemi, 44: 13-18.
- KUŚKA, A. & KANIA, I. (2010) - New soldier beetles (Coleoptera, Cantharidae) from the Eocene Baltic amber. Zootaxa, 2400: 49-56.
- KUŚKA, A. & KUPRYJANOWICZ, J. (2005) - Soldier beetles (Coleoptera: Cantharidae) from Baltic amber. Polskie Pismo Entomologiczne, 74: 309-316.
- KUZMINA, S. A. (2015) - Quaternary Insects and Environment of Northeastern Asia. Paleontological Journal, 49(7): 679-867.
- LARSSON, S. G. (1965) - Reflections on the Baltic Amber Inclusions. Entomologiske Meddelelser, 34: 135-141.
- LARSSON, S. G. (1978) - Baltic Amber – a Palaeobiological Study. Entomonograph, Vol. 1. Klampenborg, Denmark, 192 pp.
- LEE, C. M., CHAN, K. W., HO, K. H. (1997) - Palaeontology and stratigraphy of Hong Kong. Vol. 1. Beijing: Science Press, 206 p., 57 plates [in Chinese].
- LIN, Q. B. (1997) - [new taxa/taxonomic names]. In: LIN, Q. B. & LEE, C. M. (1997), Early Tertiary insects from Peng Chau, Hong Kong. Palaeontology and stratigraphy of Hong Kong, 1: 185-199, 6 plates. [in Chinese].
- LUCAS, R. (1915) - Coleoptera für 1914. Archiv für Naturgeschichte, 81: 1-227.
- MATTHEWS JR., J. V. & TELKA, A. (1997) - Insect Fossils from the Yukon (pp. 911-962). In: DANKS, H. V. & DOWNES, J. A. (Eds.), Insects of the Yukon. Biological Survey of Canada (Terrestrial Arthropods), Ottawa, 1034 pp.
- MCKENNA, D. D., WILD, A. L., KANDA, K., BELLAMY, C. L., BEUTEL, R. G., CATERINO, M. S., FARNUM, C. W., HAWKS, D. C., IVIE, M. A., JAMESON, M. L., LESCHEN, R. A. B., MARVALDI, A. E., MCHUGH, J. V., NEWTON, A. F., ROBERTSON, J. A., THAYER, M. K., WHITING, M. F., LAWRENCE, J. F., ŚLIPIŃSKI, A., MADDISON, D. R., FARRELL, B. D. (2015) - The beetle tree of life reveals that Coleoptera survived end-Permian mass extinction to diversify during the Cretaceous terrestrial revolution. Systematic Entomology, 40: 835-880.
- MENGE, F. A. (1856) - Lebenszeichen vorweltlicher, im bernstein eingeschlossener thiere. Programm, womit zu der am Montag den 17. März 1856 von 8 ½ Uhr Vorm. und 2 ½ Uhr Nachm. an stattfindenden öffentlichen

- Prüfung der Schüler der Petrischule. Danzig, A. W. Kafemann, 32 pp.
- MERRITT, J. W., AUTON, C. A., CONNELL, E. R., HALL, A. M. and PEACOCK, J. D. (2003) - Cainozoic geology and landscape evolution of north-east Scotland. British Geological Survey, Edinburgh, 178 pp., 52 figs., 27 plates, 11 maps, CD-ROM.
- MEUNIER, F. A. (1915) - Über einige fossile Insekten aus den Braunkohlenschichten (Aquitaniens) von Rott im Siebengebirge. 3. Teil. Zeitschrift der Deutschen Geologischen Gesellschaft (A), 67: 219-230, t. XXVI-XXVII.
- MEUNIER, F. A. (1920a) - Paléozoologie: Insectes. [review article of:] Wickham. H.-F., New Miocene Coleoptera from Florissant. Mus. Compar. Zoology, vol. LVIII, pp. 423-494, pl. 1-16. Cambridge, 1914. Revue critique de Paléozoologie et de Paléophytologie,
- MEUNIER, F. A. (1920b) - Paléozoologie: Insectes. [review article of:] Wickham. H.-F., The Princeton Collection of Fossil Beetles from Florissant. Ann. Entomological Soc. of America, vol. VI, pp. 359-366, pl. 38-41. Iowa. Revue critique de Paléozoologie et de Paléophytologie, 24 (4): 159.
- MIGHALL, T. M., TIMBERLAKE, S., CLARK, S. H. E., CASELDINE, A. E. (2002) - A Palaeoenvironmental Investigation of Sediments from the Prehistoric Mine of Copa Hill, Cwmystwyth, mid-Wales. Journal of Archaeological Science, 29: 1161-1188.
- MISKIMEN, G. W. (1961) - Zoogeography of the Coleopterous Family Chauliognathidae. Systematic Zoology, 10(3): 140-153.
- MITCHELL, A. A. (2013) EDNA, The Fossil Insect Database - <http://edna.palass-hosting.org> [last visite: December 2016].
- MOTSCHULSKY, V. de (1857) - Voyages. Lettres de M. de Motschulsky à M. Ménétriés. N. 4. Études Entomologiques [1856], 5
- NICHOLSON, D. B (2012) - Fossil perspectives on the evolution of insect diversity. PhD thesis, University of York, York, UK, 421
- NUDDS, J. R & SELDEN, P. A. (2008) - Fossil Ecosystems of North America. A guide to the sites and their extraordinary biotas. Manson Publishing, CRC Press, 288 pp.
- OKTAR, A. / [HARUN YAHYA] (2006-2008) - Atlas of Creation. Volume 1. Global Publishing, Istanbul, Turkey, 904 pp.
- OKUSHIMA, Y. (2005) - A Taxonomic Study on the genus Lycocerus (Coleoptera, Cantharidae) from Japan, with Zoogeographical Considerations. Japanese Journal of Systematic Entomology (Matsuyama), Monographic Series, 2: 1-383.
- OUSTALET, J.-F. É. (1874) - Recherches sur les insectes fossiles des terrains tertiaires de la France. G. Masson Éditeur, Paris (Thèses présentées à la faculté des sciences de Paris), 1-556, pl. 1-12.
- PANOVA, N. K., JANKOVSKA, V., KORONA, O. M. and ZINOV'EV, E. V. (2003) - The Holocene Dynamics of Vegetation and Ecological Conditions in the Polar Urals. Ekologiya, 4: 248-260 [in Russian. English translation: Russian Journal of Ecology, 34(4): 219-230].
- PAWŁOWSKI, J. S. & MAZUR, M. A. (2012) - Antoni Kuśka (*1940 †2010) – entomolog, pedagog, alpinista. Biografia, spis publikacji oraz wspomnienia o Zmarłym przedstawicieli dwóch pokoleń koleopterologów. Wiadomości Entomologiczne (Sylwetki Entomologów. Entomologists), 31(1): 41-56. [in Polish].
- PELLETIER, G., HÉBERT, C. (2014) - The Cantharidae of Eastern Canada and Northeastern United States. Canadian Journal of Arthropod Identification, 25 (Febbraio, 2014): 1-246. (online version, doi: 10.3752/cjai.2014.25).
- PENNEY, D. & JEPSON, J. E. [with artwork by Richard Bizley] (2014) - Fossil Insects: An introduction to palaeoentomology. Siri Scientific Press, 224 pp.
- PERIS, D., RUZZIER, E., PERRICHOT, V., DELCLÒS, X. (2016a) - Evolutionary and paleobiological implications of Coleoptera (Insecta) from Tethyan-influenced Cretaceous ambers. Geoscience Frontiers, 7: 695-706. (doi: 10.1016/j.gsf.2015.12.007 [available online 16 January 2016]).
- PERIS, D., RUZZIER, E., PERRICHOT, V., PEÑALVER, E., DELCLÒS, X. (2016b) - Paleobiology of beetles (Insecta: Coleoptera) from Western European Cretaceous ambers. 7th International Conference on Fossil Insects, Arthropods and Amber, (Poster - April 2016), Edinburgh, UK, doi: 10.13140/RG.2.1.3731.0328
- PERIS, D., SÁNCHEZ-GARCÍA, A., SORIANO, C. & DELCLÒS, X. (2013) - Beetle fauna in the Early Cretaceous Spanish amber (pp. 74-75). The 6th International Congress on Fossil Insects, Arthropods and

- Amber, Abstract Book, Byblos, Lebanon, April 14-18, 2013 [Poster].
- PERKOVSKY, E. E. (2015) - Toponyms and ethnonyms in the names of Rovno amber animals and plants. *Vestnik zoologii*, 49(5):
- PERKOVSKY, E. E., ZOSIMOVICH, V. Y., VLASKIN, A. P. (2010) - Rovno Amber (pp. 117-137). In: PENNEY, D. (Ed.). *Biodiversity of fossils in Amber from the major world deposits*. Siri Scientific Press, Manchester, 304 pp.
- PERRILLIAT, M. D. C. (1989) - Insecta (pp. 388-397). In: CARREÑO, A. L., PERRILLIAT, M. D. C., GONZÁLEZ-ARREOLA, C., APPLEGATE, S. P., CARRANZA-CASTAÑEDA, O., MARTÍNEZ-HERNÁNDEZ, E. (eds.), *Fósiles Tipo Mexicanos*. México, D. F., Universidad Nacional Autónoma de México. Instituto de Geología, 531 pp.
- PETRULEVICIUS, J. F. (1999) - Insectos del Cenozoico de la Argentina. *Revista de la Sociedad Entomológica Argentina*, 58(1-2): 95-103.
- PICTET, F. J. (1846) - *Traité élémentaire de Paléontologie ou histoire naturelle des animaux fossiles considérés dans leurs rapports zoologiques et géologiques*. Tome Quatrième. Imprimerie de Jules-Guillaume Fick., Genève, xiv, + 1, 458 pp, 20 pls.
- PICTET, F. J. (1854) - *Traité de Paléontologie ou histoire naturelle des animaux fossiles considérés dans leurs rapports zoologiques et géologiques* (Seconde Édition, revue, corrigée, considérablement augmentée, accompagnée d'un atlas de 110 planches grand in-4°). Tome Deuxième. J.-B. Baillièvre, Paris, 727 pp.
- PIELIŃSKA, A. (2006) - XXI Spotkanie Badaczy Bursztynu W Muzeum Ziemi PAN, Warszawa, 5 grudnia 2006 roku (pp. 8-10). The 21st Amber Researchers' Meeting at the Museum of the Earth, Polish Academy of Sciences (PAN), Warsaw, 5 December 2006 (pp. 32-34). Bursztynisko, 27 [bilingual: in Polish and English].
- PITON, L. (1939) - Note complémentaire sur les insectes fossiles des cinérites pliocènes du Lac Chambon (Puy-de-Dôme). *Revue des Sciences Naturelles d'Auvergne* (N. S.), 5: 102-108.
- PITON, L. & THÉOBALD, N. (1936) - Les Insectes fossiles du Puy-de-Mur (Limagne). *Bulletin Mensuel de la Société des Sciences de Nancy*, 1936, (N. S.), 9: 202-214.
- POINAR JR., G. O. (1992) - *Life in Amber*. Stanford University Press, Stanford, 368 pp.
- POINAR JR., G. O. & FANTI, F. (2016) - New fossil soldier beetles (Coleoptera: Cantharidae) in Burmese, Baltic and Dominican amber. *Palaeodiversity*, 9: 1-7; Stuttgart 30 December 2016. Published online: 31 May 2016, DOI: 10.18476/pale.v9.a1
- POINAR Jr., G. O., MARSHALL, C. J., BUCKLEY, R. (2007) - One Hundred Million Years of Chemical Warfare by Insects. *Journal of Chemical Ecology*, 33(9): 1663-1669.
- PONOMARENKO, A. G. (1992) - Upper Liassic beetles (Coleoptera) from Lower Saxony, Germany. *Senckenbergiana lethaea*, 72: 179-188.
- PONOMARENKO, A. G. (1995) - The geological history of beetles (pp. 155-171). In: PAKALUK, J. & ŚLIPIŃSKI, S. A. (eds.), *Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson*. Vol. 1. Muzeum i Instytut Zoologii PAN, Warsaw.
- PONOMARENKO, A. G. & SCHULTZ, O. (1988) - Typen der Geologisch-Paläontologischen Abteilung: Fossile Insekten. Kataloge der wissenschaftlichen Sammlung des Naturhistorischen Museums in Wien, 6 (Paläozool. 1): 39 pp. [+ 1], 14 tables.
- RAMSDALE, A. S. (2002) - Family 64. Cantharidae Imhoff 1856 (pp. 202-218). In: ARNETT, R. H., Jr., THOMAS, M. C., SKELLEY, P. E. and FRANK, J. H. (Eds.) *American Beetles*, Volume 2. Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 880 pp.
- RASNITSYN, A. P., BASHKUEV, A. S., KOPYLOV, D. S., LUKASHEVICH, E. D., PONOMARENKO, A. G., POPOV, Y. A., RASNITSYN, D. A., RYZHKOVA, O. V., SIDORCHUK, E. A., SUKATSHEVA, I. D., VORONTSOV, D. D. (2016) - Sequence and scale of changes in the terrestrial biota during the Cretaceous (based on materials from fossil resins). *Cretaceous Research*, 61: 234-255 + Appendix A and Appendix B.
- RASNITSYN, A. P. & ROSS, A. J. (2000) - A preliminary list of arthropod families present in the Burmese amber collection at The Natural History Museum, London. *Bulletin of the Natural History Museum, London (Geology Series)*, 56(1): 21-24.
- ROHDENDORF, B. B., PONOMARENKO, A. G. (1962) - Otryad Coleoptera. Žhestkokriliye, ili Žuki. *Osnovy Paleontologii*, Tom. 9: Chlenistonogie Trakheinie i Khelitserovie (Ed. B.B. Rohdendorf), Akademia Nauk SSR, Moscow, pp. 241-267 [in Russian]. [English translation: ROHDENDORF, B. B., PONOMARENKO, A. G. (1991) - Order Coleoptera. Beetles. Fundamentals of Paleontology, Volume 9. Arthropoda, Tracheata,

- Chelicerata (Ed. B.B. Rohdendorf), Smithsonian Institution Libraries and The National Science Foundation, Washington, D.C., 341-382].
- ROSS, A. J., MELLISH, C., YORK, P., CRIGHTON, B. (2010) - Burmese Amber (pp. 209-236). In: PENNEY, D. (Ed.). Biodiversity of fossils in Amber from the major world deposits. Siri Scientific Press, Manchester, 304 pp.
- SCHAUFUSS, C. F. C. (1892) - Preussens Bernstein-Käfer. Neue Formen aus der Helm'schen Sammlung im Danziger Provinzialmuseum. Berliner Entomologische Zeitschrift, 36(I) [1891]: 53-64.
- SCHAWALLER, W. (1986) - Fossile Käfer aus miozänen Sedimenten des Randecker Maars in Südwest-Deutschland (Insecta: Coleoptera). Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie), 126: 1-9.
- SCHRÖDER, C. (1900) - Litteratur-Referate. [review article of:] Helm, Otto: Tierische Einschlüsse in Succinit. 3 p. In: "Schrift. Naturf. Ges. Danzig", N. F., Bd. IX, 2. Illustrierte Zeitschrift für Entomologie, 5: 355.
- SCOTT, R., SANTUCCI, V. L. and CONNORS, T. (2001) - An inventory of paleontological resources from the national parks and monuments in Colorado (pp. 178-202). In: SANTUCCI V. L. & MCCLELLAND L. (eds.), Proceedings of the 6TH fossil resource Conference, Geologic Resources Division Technical Report.
- SCUDDER, S. H. (1876) - Fossil Coleoptera from the Rocky Mountain Tertiaries. Bulletin of the United States Geological and Geographical Survey of the Territories, 2(I): 77-87.
- SCUDDER, S. H. (1882) - The Tertiary Lake-basin at Florissant, Colorado, between South and Hayden Parks. Bulletin of the United States Geological and Geographical Survey of the Territories, 6: 279-300.
- SCUDDER, S. H. (1885) - Systematische Übersicht der Fossilen Myriopoden, Arachnoideen und Insekten. In: ZITTEL, K. A. (Ed.), Handbuch der Palaeontologie. I. Abtheilung. Palaeozoologie. Bd. 2. München und Leipzig, R. Oldenbourg, 721-831. [French translation: SCUDDER, S. H. (1887)].
- SCUDDER, S. H. (1886) - Systematic review of our present knowledge of fossil insects including myriapods and arachnids. Bulletin of the United States Geological Survey, 31: 1-128.
- SCUDDER, S. H. (1887) - Myriapodes, Arachnides, Insectes. In: ZITTEL, K. A. (Ed.), Traité de Paléontologie. Tome II. Paléozoologie, Partie I: Mollusca et Arthropoda. Paris: Doin et Oldenbourg, 720-833. [original version in German: SCUDDER, S. H. (1885)].
- SCUDDER, S. H. (1891) - Index to the known fossil insects of the world including myriapods and arachnids. Bulletin of the United States Geological Survey, 71: 1-744.
- SCUDDER, S. H. (1895) - The Miocene Insect-fauna of Oeningen, Baden. The Geological Magazine: or, Monthly Journal of Geology, n. s., (Decade IV), 2: 116-122 + 1 plate.
- SCUDDER, S. H. (1900) - Adephagous and Clavicorn Coleoptera from the Tertiary deposits at Florissant, Colorado with descriptions of a few other forms and a systematic list of the non-rhynchophorous Tertiary Coleoptera of North America. Monographs of the United States Geological Survey, 40: 1-148, 11 pls.
- SENDEL, N. (1742) - Historia succinorum corpora aliena involventium et naturae opere pictorum et caelatorum ex AUGUSTORUM I et II cimeliis [or "ex REGIIS AUGUSTORUM cimeliis"] Dresden conditis aeri insculptorum. Gleditsch, Leipzig, 328 pp, 13 t. [title with two versions].
- SERRES, P. M. T. de (1843) - Notes géologiques sur la Provence. Actes de la Société Linnéenne de Bordeaux, 13: 1-82.
- SHORT, S. K., ELIAS, S. A., WAYTHOMAS, C. F. and WILLIAMS, N. E. (1992) - Fossil Pollen and Insect Evidence for Postglacial Environmental Conditions, Nushagak and Holitna Lowland Regions, Southwest Alaska. Arctic, 45(4): 381-392.
- SMITH, D. N., ROSEFF, R., BEVAN, L., BROWN, A. G., BUTLER, S., HUGHES, G. and MONCKTON, A. (2005) - Archaeological and environmental investigations of a Lateglacial and Holocene river sedimentary sequence on the River Soar at Croft, Leicestershire, UK. - The Holocene, 15(3): 353-377.
- SOLÓRZANO KRAEMER, M. M. (2007) - Systematic, palaeoecology, and palaeobiogeography of the insect fauna from Mexican amber. Palaeontographica Abteilung A, 282(1-6): 1-133.
- SOLÓRZANO KRAEMER, M. M. (2010) - Mexican Amber (pp. 42-57). In: PENNEY, D. (Ed.). Biodiversity of fossils in Amber from the major world deposits. Siri Scientific Press, Manchester, 304 pp.
- SPAHR, U. (1981a) - Bibliographie der Bernstein- und Kopal-Käfer (Coleoptera). Bibliography of Coleoptera in amber and copal. Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie), 72: 21 pp.
- SPAHR, U. (1981b) - Systematischer Katalog der Bernstein- und Kopal-Käfer (Coleoptera). Systematic

- Catalogue of Coleoptera in Amber and Copal. Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie), 80: 107 pp.
- ŠVIHLA, V. (2008) - Redefinition of the subgenera of the genus *Themus* Motschulsky, 1858, with description of five new species (Coleoptera: Cantharidae). *Vernate*, 27: 183-190.
- TAKAHASHI, K. (2007) - Taxonomic Status of *Malthacus* (Coleoptera, Cantharidae). *Japanese Journal of Systematic Entomology*, 13(2): 241-251.
- THÉOBALD, N. (1937) - Les insectes fossiles des terrains oligocènes de France. Mémoires de la Société des Sciences de Nancy, Année 1937: 473 pp., 29 pls., + 1 p. [also printed as Thèse, Docteur ès Sciences Naturelles, Université de Nancy].
- THE PALEOBIOLOGY DATABASE - <https://paleobiodb.org> [last vist: December 2016].
- TOUSSAINT, E. F. A., SEIDEL, M., ARRIAGA-VARELA, E., HÁJEK, J., KRÁL, D., SEKERKA, L., SHORT, A. E. Z. & FIKÁČEK, M. (2016) - The peril of dating beetles. *Systematic Entomology*, doi: 10.1111/syen.12198
- VITALI, F. (2009) - The cerambycids included in Baltic amber: current knowledge status with the description of new taxa (Coleoptera, Cerambycidae). *Denisia*, 26, zugleich Kataloge der oberösterreichischen Landesmuseen Neue Serie 86 (2009): 231-242.
- VOLLMAR, von [...] (1835) - Ueber fossile Entomologie. *Faunus, Zeitschrift für Zoologie und vergleichende Anatomie*, 2: 56-62.
- WEDMANN, S., POSCHMANN, M., HÖRNSCHEMEYER, T. (2010) - Fossil insects from the Late Oligocene Enspel Lagerstätte and their palaeobiogeographic and palaeoclimatic significance. *Palaeobiodiversity and Palaeoenvironments*, 90(1): 49-58.
- WEIDNER, H. (1952) - Insektenleben im Bernsteinwald. (Ein Bericht über die Bernsteinsammlung des Geologischen Staatsinstitutes Hamburg). *Entomologische Zeitschrift* [1952-1953], 62(8): 62-64 and 62(9): 65-72 and 62(11): 88.
- WEITSCHAT, W. (2009) - Jäger, Gejagte, Parasiten und Blinde Passagiere – Momentaufnahmen aus dem Bernsteinwald. *Denisia*, 26, zugleich Kataloge der oberösterreichischen Landesmuseen Neue Serie 86 (2009): 243-256.
- WEITSCHAT, W. & WICHARD, W. (1998) - Atlas der Pflanzen und Tiere im Baltischen Bernstein. Verlag: Dr. Friedrich Pfeil, 256 pp. [English translation: WEITSCHAT, W. & WICHARD, W. (2002)].
- WEITSCHAT, W. & WICHARD, W. (2002) - *Atlas of Plants and Animals in Baltic Amber*. Verlag: Dr. Friedrich Pfeil, 256 pp. [original version in German: WEITSCHAT, W. & WICHARD, W. (1998)].
- WEITSCHAT, W. & WICHARD, W. (2010) - Baltic Amber (pp. 81-115). In: PENNEY, D. (Ed.). *Biodiversity of fossils in Amber from the major world deposits*. Siri Scientific Press, Manchester, 304 pp.
- WESTWOOD, J. O. (1854) - Contributions to Fossil Entomology. *The Quarterly Journal of the Geological Society of London*, 10
- WHEELER, W. M. (1906) - The expedition to Colorado for fossil insects. *The American Museum Journal*, 6(4): 199-203.
- WHITE, R. D. (1995) - A Type Catalog of Fossil Invertebrates (Arthropoda: Hexapoda) in the Yale Peabody Museum. *Postilla*, 209: 55
- WICHARD, W. (2009) - Taphozönosen im Baltischen Bernstein. *Denisia*, 26, zugleich Kataloge der oberösterreichischen Landesmuseen Neue Serie 86 (2009): 257-266.
- WICHARD, W. & GREVEN, H. (2009) - Über Tausendfüßler, spanische Fliegen und Heuschrecken – Zur “Historia Succinorum” des Nathanael Sendel von 1742. *Denisia*, 26, zugleich Kataloge der oberösterreichischen Landesmuseen Neue Serie 86 (2009): 267-294.
- WICHARD, W. & WEITSCHAT, W. (2004) - Im Bernsteinwald. Gerstenberg Verlag, Hildesheim, 168 pp.
- WICKHAM, H. F. (1909) - New Fossil Coleoptera from Florissant. *The American Journal of Science*, (Fourth Series), 28 [Whole number, CLXXVIII]: 126-130.
- WICKHAM, H. F. (1913a) - Fossil Coleoptera from Florissant in the United States National Museum. *Proceedings of the United States National Museum*, 45: 283-303, 5 plates.
- WICKHAM, H. F. (1914) - New Miocene Coleoptera from Florissant. *Bulletin of the Museum of Comparative Zoölogy at Harvard College*, in Cambridge, 58(11): 423-494 + 1 + 16 plates (and explanation).
- WICKHAM, H. F. (1917) - New species of fossil beetles from Florissant, Colorado. *Proceedings of the United*

FOSSILS & MINERALS

States National Museum, 52: 463-472, 3 plates.

WICKHAM, H. F. (1920) - Catalogue of the North American Coleoptera described as fossils (pp. 347-365). In: LENG, C. W. (ed.), Catalogue of the Coleoptera of America, North of Mexico. John D. Sherman, Jr., New York.

WILLEMSTEIN, S. C. (1987) - An evolutionary basis for pollination ecology. E. J. Brill, Leiden University Press, Leiden, 425 pp.

WILLIAMSON, G. C. (1932) - The Book of Amber. Ernest Benn Limited, London, xiii + [i] + 15-268 pp., + 1 pl.

WILSON, M. V. H. (1978) - Paleogene insect faunas of Western North America. *Quaestiones Entomologicae*, 14: 13-34.

WILSON, M. V. H. (2008-2009) - McAbee Fossil Site Assessment, Final Report July 30, 2007. Revised August 5, 2007. Further revised October 24, 2008. Unpublished report, British Columbia Ministry of Agriculture and Lands, 60 pp.

WINKLER, J. R. (1987) - Berendtimiridae fam. n., a new family of fossil beetles from Baltic Amber. (Coleoptera, Cantharoidea). *Mitteilungen der Münchener Entomologischen Gesellschaft*, 77: 51-59.

WITTMER, W. (1963) - A new cantharid from the Chiapas amber of Mexico. *University of California Publications in Entomology*, 31: 53 + plate 3 (p. 59).

YABLOKOV-KHNZORIAN, S. M. (1960) - New beetles from the Baltic amber. *Paleontologicheskii Zhurnal*, 3: 90-101. [in Russian. Title: *Novye žhestkokrylye iz Baltiyskogo yantarya*].

YANG, Y. X. & YANG, X. K. (2010) - A redescription of the genus *Cyrebion* Fairmaire, 1891, with notes on related taxa and distribution (Coleoptera: Cantharidae). *Journal of Natural History*, 44(9-12): 579-588.

ZANG, R. (1905) - Coleoptera Longicornia aus der Berendtschen Bernsteinsammlung. *Sitzungs-Berichte der Gesellschaft Naturforschender Freunde zu Berlin*, 1905: 232-245, 1 t.

ZHANG, J. (1989) - Fossil Insects from Shanwang, Shandong, China. Shandong Science and Technology Publishing House. Jinan, China, 459 pp., 92 plates [in Chinese with English summary].

ZHANG, J., SUN, B. & ZHANG, X. (1994) - Miocene insects and spiders from Shanwang, Shandong. Science Press, Beijing, v + 298 pp., 44 plates [in Chinese with English summary].

MUSEO DEL CAVATORE

Via 17 Agosto 1944, 10/a Vellano (PT) tel. e fax 0572 409181 +39330910517



GeoElsa

MOSTRA DI MINERALI - FOSSILI CONCHIGLIE

8 - 9 Aprile 2017

**Palazzetto
dello
Sport**

Via Liguria, 1

**Colle
di
Val d'Elsa (SI)**

Ore 09:00 - 19:00

Informazioni:

info@geoelsa.it
www.geoelsa.it

Organizzazione:

Associazione Mineralogica
e Paleontologica Senese

Gruppo Mineralogico Senese

Gruppo Paleontologico "C. De Giuli"



Comune di
Colle di Val d'Elsa



INGRESSO LIBERO